

WEST[Help](#)[Logout](#)[Interrupt](#)[Main Menu](#)[Search Form](#)[Posting Counts](#)[Show S Numbers](#)[Edit S Numbers](#)[Preferences](#)**Search Results -**

Terms	Documents
(116 or 115 or 114 or 113 or 112 or 111 or 110) and 18 and 16 and 11	3

Database:

US Patents Full-Text Database
US Pre-Grant Publication Full-Text Database
JPO Abstracts Database
EPO Abstracts Database
Derwent World Patents Index
IBM Technical Disclosure Bulletins

Refine Search:

(116 or 115 or 114 or 113 or 112 or 111
or 110) and 18 and 16 and 11

[Clear](#)**Search History****Today's Date: 5/10/2001**

<u>DB Name</u>	<u>Query</u>	<u>Hit Count</u>	<u>Set Name</u>
USPT,PGPB	(l16 or l15 or l14 or l13 or l12 or l11 or l10) and l8 and l6 and l1	3	<u>L17</u>
USPT,PGPB	(((435/938)!.CCLS.))	100	<u>L16</u>
USPT,PGPB	(((435/921)!.CCLS.))	308	<u>L15</u>
USPT,PGPB	(((435/320.1)!.CCLS.))	9069	<u>L14</u>
USPT,PGPB	(((435/254.23)!.CCLS.))	86	<u>L13</u>
USPT,PGPB	(((435/254.22)!.CCLS.))	52	<u>L12</u>
USPT,PGPB	(((435/142)!.CCLS.))	93	<u>L11</u>
USPT,PGPB	((435/136)!.CCLS.)	329	<u>L10</u>
USPT,PGPB	l8 and l1 and l2 and l6	2	<u>L9</u>
USPT,PGPB	prepar\$7 or synthes\$4 or mak\$4	1756877	<u>L8</u>
USPT,PGPB	l6 and l4	2	<u>L7</u>
USPT,PGPB	Hydroxylation and alkane\$1	389	<u>L6</u>
USPT,PGPB	Hydroxylation or alkane\$1	33596	<u>L5</u>
USPT,PGPB	l2 and l1	52	<u>L4</u>
USPT,PGPB	pichia pastoris	972	<u>L3</u>
USPT,PGPB	candida maltosa or Candida cloacae or Candida novellus or Candida subtropicalis	177	<u>L2</u>
USPT,PGPB	dicarboxylic acid\$1 or Carboxylic acid\$1 or monocarboxylic acid\$1	155036	<u>L1</u>

WEST**Generate Collection****Search Results - Record(s) 1 through 2 of 2 returned.**☐ 1. Document ID: US 6020288 A

L7: Entry 1 of 2

File: USPT

Feb 1, 2000

US-PAT-NO: 6020288

DOCUMENT-IDENTIFIER: US 6020288 A

TITLE: Methods and compositions for enhancing cytochrome P450 in plants

DATE-ISSUED: February 1, 2000

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Nonomura; Arthur M.	Boxborough	MA	01719	N/A
Benson; Andrew A.	La Jolla	CA	92037	N/A
Nishio; John N.	Laramie	WY	82070-3917	N/A

APPL-NO: 8/ 927415

DATE FILED: September 11, 1997

PARENT-CASE:

RELATED APPLICATIONS This application is a continuation-in-part of U.S. patent application Ser. No. 08/610,928, filed Mar. 5, 1996, now U.S. Pat. No. 5,846,908, which is a continuation-in part of U.S. patent application Ser. No. 08/399,399, filed Mar. 6, 1995; which was a continuation-in part of Ser. No. 08/351,348 filed Dec. 9, 1994, now U.S. Pat. No. 5,597,400 issued on Jan. 28, 1997, which was a continuation-in-part of U.S. patent application Ser. No. 07/901,366, filed on Jun. 19, 1992. The full disclosures of each of these patent applications are incorporated herein by reference. Related international application are PCT/US96/02444 (equivalent of Ser. No. 08/610,928) and PCT/US93/05676 (equivalent of Ser. No. 08/351,348).

INT-CL: [6] A01N 31/00, A01N 37/00, A01N 43/22, A01N 57/02

US-CL-ISSUED: 504/127; 504/128, 504/130, 504/136, 504/138, 504/140, 504/142, 504/143, 504/144, 504/149

US-CL-CURRENT: 504/127; 504/128, 504/130, 504/136, 504/138, 504/140, 504/142, 504/143, 504/144, 504/149

FIELD-OF-SEARCH: 504/127, 504/128, 504/130, 504/136, 504/138, 504/140, 504/142, 504/143, 504/144, 504/149

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<u>3897241</u>	July 1975	Washio et al.	71/113
<u>4799953</u>	January 1989	Danzig et al.	71/98
<u>4846877</u>	July 1989	Azuma et al.	71/92
<u>5298482</u>	March 1994	Tanaka et al.	504/320
<u>5300540</u>	April 1994	Masters	523/309
<u>5532204</u>	July 1996	Joshi	504/118
<u>5597400</u>	January 1997	Nonomura et al.	71/28

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
0 465 907 A1	January 1992	EPX	
2689905	October 1993	FRX	
2004856	April 1979	GBX	

OTHER PUBLICATIONS

Whitlock, J.P. et al. Induction of Cytochrome P450 Enzymes That Metabolize Xenobiotics. Chapter 10 in Cytochrome P450: Structure, Mechanism and Biochemistry, 2nd ed. Ortiz de Montellano, ed. 1995, pp. 367-389.

Porter, T.D. et al. Chytochrome P-450. The Journal of Biological Chemistry. Jul. 25, 1991, vol. 266, No. 21 pp. 13469-13472.

Bollwell, G.P. et al. Plant Cytochrome P450. Phytochemistry. 1994, vol. 37, No. 6, pp. 1491-1506.

Schuler, M.A. Plant Cytochrome P450 Monooxygenases. Critical Reviews in Plant Sciences. 1996, vol. 15, No. 3, pp. 235-284.

Halkier, B.A. et al. Involvement o Cytochrome P-450 in the Biosynthesis of Dhurrin in Sorghum bicolor (L.) Moench. Plant Physiology. 1991, vol. 96, pp. 10-17.

Aharoni, A. et al., "Isolation and Characterization of Cytochrome P-450 cDNAs from Strawberry Fruit" 17th Intl. Congress of Biochem. and Molec. Biol., San Francisco, CA, p. A811, Abstract No. P233 (Aug. 24-29, 1997).

Angerhorfer, A. and Bittl, R., "Radicals and Radical Pairs in Photosynthesis" Photochem. Photobiol. 63(1):11-38 (1996).

Badger, M.R. and Schrieber, U., "Effects of inorganic carbon accumulation on photosynthetic oxygen reduction and cyclic electron flow in the cyanobacterium Synechococcus PCC7942" Photosynthesis Res. 37:177-191 (1993).

Bolwell, G.P. et al., "Plant Cytochrome P450" Phytochemistry 37(6):1491-1506 (1994).

Bowling, S.A. et al., "A mutation in Arabidopsis that leads to constitutive Expression of systemic acquired resistance" The Plant Cell 6:1845-1857 (Dec. 1994).

Butler, J. and Hoey, B.M., "The one-electron reduction potential of several substrates can be related to their reduction rates by cytochrome P-450 reductase" Biochimica et Biophysica Acta 1161:73-78 (1993).

Nee, M.W. and Bruice, T.C., "Use of the N-Oxide of p-Cyano-N,N-dimethylaniline as an "Oxygen" Donor in a Cytochrome P-450 Model System" J. Am. Chem. Soc. 104(22):6123-6125 (1982).

Nonomura, A.M. and Benson, A.A., "The path of carbon in photosynthesis: Improved crop yields with methanol" Proc. Natl. Acad. Sci. USA 89:9794-9798 (1992).

Ohkawa, H. et al., "Genetically Engineered Plants Expressing Mammalian P450 Monooxygenases for Phytoremediation" 17th Intl. Congress of Biochem. and Molec. Biol., San Francisco, CA, Abstract No. P37 (Aug. 24-29, 1997).

Palazon, J. et al., "Effects of Auxin and Phenobarbital on Morphogenesis and Production of Digitoxin in Digitalis callus" Plant Cell Physiol. 36(2):247-252 (1995).

Parikh, A. et al., "Drug metabolism by Escherichia coli expressing human

- cytochromes P450" *Nature Biotechnol.* 15:784-788 (1997).
- Porter, T.D. and Coon, M.J., "Cytochrome P-450" *J. Biol. Chem.* 266:13469-13472 (1991).
- Sasame, H.A. and Gillette, J.R., "Studies on the Relationship between the Effects of Various Substances on Absorption Spectrum of Cytochrome P-450 and the Reduction of p-Nitrobenzoate by Mouse Liver Microsomes" *Mol. Pharmacol.* 5:123-130 (1969).
- Schuler, M.A., "Plant Cytochrome P450 Monooxygenases" *Crit. Rev. Plant Sci.* 15(3):235-284 (1996).
- Sherry, B. and Abeles, R.H., "Mechanism of Action of Methanol Oxidase, Reconstitution of Methanol Oxidase with 5-Deazaflavin, and Inactivation of Methanol Oxidase by Cyclopropanol" *Biochem.* 24(11):2594-2605 (1985).
- Strobel, H.W. et al., "NADPH Cytochrome P450 Reductase and Its Structural and Functional Domains" in: *Cytochrome P450, Structure, Mechanism, and Biochemistry*, Second Edition, Paul R. Ortiz de Montellano, Ed., Plenum Press, NY, pp. 225-390 (1995).
- Szekeres, M. et al., "Brassinosteroids Rescue the Deficiency of CYP90, a Cytochrome P450, Controlling Cell Elongation and De-etiolation in Arabidopsis" *Cell* 85:171-182 (1996).
- Takabe, T., "Glycolate Formation Catalyzed by Spinach Leaf Transketolase Utilizing the Superoxide Radical" *Biochem.* 19:3985-3989 (1980).
- Tolbert, N.E. et al., "The oxygen and carbon dioxide compensation points of C.sub.3 plants: Possible role in regulating atmospheric oxygen" *Proc. Natl. Acad. Sci. USA* 92:11230-11233 (1995).
- Wachtveitl, J. et al., "Tyrosine 162 of the Photosynthetic Reaction Center L-Subunit Plays a Critical Role in the Cytochrome C.sub.2 Mediated Rereduction of the Photooxidized Bacteriochlorophyll Dimer in Rhodobacter sphaeroides" *Biochem.* 32:10894-10904 (1993).
- Wardman, P., "Reduction Potentials of One-Electron Couples Involving Free Radicals in Aqueous Solution" *J. Phys. Chem. Ref. Data* 18(4):1637-1755 (1989).
- Wendler, C. et al., "Effect of Glufosinate (Phosphinothricin) and Inhibitors of Photorespiration on Photosynthesis and Ribulose-1,5-Bisphosphate Carboxylase Activity" *J. Plant Physiol.* 139:666-671 (1992).
- Cottrell, S. et al., "Studies on the cytochrome P-450 of avocado (*Persa americana*) mesocarp microsomal fraction" *Xenobiotica* 20(7):711-726 (1990).
- Damme, B. et al., "Induction of hepatic cytochrome P4502E1 in rats by acetylsalicylic acid or sodium salicylate" *Toxicology* 106:99-103 (1996).
- Frey, M. et al., "Analysis of a Chemical Plant Defense Mechanism in Grasses" *Science* 277:696-699 (Aug. 1, 1997).
- Gruber, V. et al., "Human haemoglobin from transgenic tobacco" *Nature* 386:29-30 (Mar. 6, 1997).
- Halkier, B.A. and Moller, B.L., "Involvement of Cytochrome P-450 in the Biosynthesis of Dhurrin in *Sorghum bicolor* (L.) Moench" *Plant Physiol.* 96:10-17 (1991).
- Hara, Y. et al., "Effect of Gibberellic Acid on Berberine and Tyrosine Accumulation in *Coptis japonica*" *Photochem.* 36(3):643-646 (1994).
- He, S. et al., "The surface-exposed tyrosine residue Tyr83 of pea plastocyanin is involved in both binding and electron transfer reactions with cytochrome f" *EMBO J.* 10(13):4011-4016 (1991).
- Holmberg, N. et al., "Transgenic tobacco expressing *Vitreoscilla* hemoglobin exhibits enhanced growth and altered metabolite production" *Nature Biotechnol.* 15:244-247 (Mar. 15, 1997).
- Kargel, E. et al., "*Candida maltosa* NADPH-cytochrome P450 Reductase: Cloning of a Full-length cDNA, Heterologous Expression in *Saccharomyces cerevisiae* and Function of the N-terminal Region for Membrane Anchoring and Proliferation of the Endoplasmic Reticulum" *Yeast* 12:333-348 (1996).
- Klein, M.L. and Fulco, A.J., "Critical residues involved in FMN binding and catalytic activity in Cytochrome P450.sub.BM-3" *J. Biol. Chem.* 268(10):7553-7561 (1993).
- Koch, B.M. et al., "The Primary Sequence of Cytochrome P450_{tyr}, the Multifunctional N-Hydroxylase Catalyzing the Conversion of L-Tyrosine to p-Hydroxyphenylacetaldehyde Oxime in the Biosynthesis of the Cyanogenic Glucoside Dhurrin in *Sorghum bicolor* (L.) Moench" *Archives Biochem. Biophys.* 323(1):177-186 (1995).
- Kusukawa, M. and Iwamura, H., "N-(3,4-Methylenedioxyphenyl) carbamates as

Potent Flower-Inducing Compounds in Asparagus Seedlings as well as Probes for Binding to Cytochrome P-450" Z. Naturforsch 50c:373-379 (1995).
 Luo, M. et al., "Characterization of a Gene Family Encoding Absciscic Acid--and Environmental Stress-inducible Proteins of Alfalfa" J. Biol. Chem. 267(22):15367-15374 (1992).
 MacDonald, G.M. et al., "A difference Fourier-transform infrared study of two redox-active tyrosine residues in photosystem II" Proc. Natl. Acad. Sci. USA 90:11024-11028 (Dec. 1993).
 Miles, C.S. et al., "Tyr-143 facilitates interdomain electron transfer in flavocytochrome b.sub.2 " Biochem. J. 285:187-192 (1992).
 Miller, A.G. and Canvin, D.T., "Glycolaldehyde Inhibits CO.sub.2 Fixation in the Cyanobacterium Synechococcus UTEX 625 without Inhibiting the Accumulation of Inorganic Carbon or the Associated Quenching of Chlorophyll a Fluorescence" Plant Physiol. 91:1044-1049 (1989).
 Moreland, D.E. et al., "Metabolism of Metolachlor by a Microsomal Fraction Isolated from Grain Sorghum (Sorghum bicolor) Shoots" Z. Naturforsch 45c:558-564 (1990).

ART-UNIT: 166

PRIMARY-EXAMINER: Clardy; S. Mark

ATTY-AGENT-FIRM: Nields, Lemack & Dingman

ABSTRACT:

The present invention provides methods for treating plants which comprise application of an oxidant that induces NADPH:cytochrome P450 reductase and application of a reductant that induces cytochrome P450 monooxygenase. The present invention also provides methods for increasing cytochrome P450 in plants and for enhancing the growth of plants. The present invention also provides compositions and systems useful in the methods of the present invention.

38 Claims, 1 Drawing figures

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWC	Draw Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-----------	-------

☐ 2. Document ID: US 5254466 A

L7: Entry 2 of 2

File: USPT

Oct 19, 1993

US-PAT-NO: 5254466

DOCUMENT-IDENTIFIER: US 5254466 A

TITLE: Site-specific modification of the candida tropicals genome

DATE-ISSUED: October 19, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Picataggio; Stephen	Santa Rosa	CA	N/A	N/A
Deanda; Kristine	Grafton	CA	N/A	N/A
Eirich; L. Dudley	Santa Rosa	CA	N/A	N/A

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Henkel Research Corporation	Santa Rosa	CA	N/A	N/A	02

APPL-NO: 7/ 432091

DATE FILED: November 6, 1989

INT-CL: [5] C12N 15/09, C12P 7/44

US-CL-ISSUED: 435/142; 435/172.1, 435/172.3, 435/924, 435/254.22

US-CL-CURRENT: 435/142; 435/254.22, 435/477, 435/481, 435/483, 435/484,
435/490, 435/6, 435/924

FIELD-OF-SEARCH: 435/67.1, 435/142, 435/172.3, 435/172.1, 435/940, 435/255,
435/924, 935/22, 935/28

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<u>4735901</u>	April 1988	Kurtz et al.	435/172.3

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
183070	April 1986	EPX	

OTHER PUBLICATIONS

Okazaki et al. Gene 57:37-44 (1987).
Okazaki et al. Biochem. 83:1232-1236 (1986).
Kelly et al. Mol. Cell. Biol. Jan 1987: p. 199-207.
Gillam et al. Mol. Gen. Genet. 198: 179 (1984).
Stromnaes et al. J. Bact. Jan. 1986: 197.
Ditchburn et al. J. Gen Microbiol. 67:299 (1971).
Gaillasdin et al. Chem Abst. vol. 77 (23) No. 149584R 1971.
Watanabe et al. Chem. Abst. vol. 70 (15) No. 65372n 1968.
Small et al. J. Cell. Biol. 105:247 (1987).
Strain and Species Identification by Restriction . . . Candida Species, Journal
of Bacteriology, Apr. 1987, pp. 1639-1643, Magee et al.
The Genetics of Candida, CRC Press, 1990, pp. 177-186, Kirsch et al.
Genetics of Candida albicans, Microbiological Reviews, Sep. 1990, pp. 226-241,
Scherer et al.
Isolation and Determination of Yeasts . . . Source of Carbon, Agr. Biol. Chem.,
vol. 30, No. 12, pp. 1175-1182, 1966, Tanabe et al.
Methods for the Genetics and Molecular Biology of Candida albicans, Analytical
Biochemistry, 175, pp. 361-372 (1988), Magee et al.
The Genus Candida Berkhout nom. conserv. . . . Delimitation, System. Appl.
Microbiol. 12, pp. 183-190 (1989), Viljoen et al.
Variation in the electrophoretic karyotype . . . in Candida albicans, Journal
of General Microbiology (1990), 136, pp. 2433-2442, Iwaguchi et al.
The Carboxyl-terminal Tripeptide Ala-Lys-Ile . . . Yeast Peroxisomes, The
Journal of Biological Chemistry, vol. 266, No. 34, pp. 23197-23208, 1991,
Aitchison et al.
In vivo import of Candida tropicalis . . . Candida albicans, Current Genetics,
1990, 17:481-486, Aitchison et al.
Redefinition of Candida Berkhout and the consequent emendation of Cryptococcus
Kuetzing and Rhodotorula Harrison, accepted 31 Mar. 1988, Weijman et al.
Interspecific Complementation Analysis . . . Candida albicans Adenine
Auxotrophs, Journal of Bacteriology, Jun. 1989, vol. 171, No. 6, pp. 3586-3589,
Corner et al.
Sequence and transcript analysis . . . -phosphate decarboxylase, Curr Genet
(1989), 16:153-157, Losberger et al.
The yeasts a taxonomic study, third edition, Elsevier Science Publishers
B.V.--Amsterdam, N.J.W. Kreger-van Rij, ed.
Differential Identification of Candida Species . . . Polypeptide Profiles,
Analytical Biochemistry 175, 548-551 (1988), Shen et al.

Transformation of Intact Yeast Cells Treated with Alkali Cations, Hisao Ito, Yasuki Fukuda, Kousaku Murata and Akira Kimura, Jan. 1983, pp. 163-168.

Acyl-CoA Oxidase From *Candida tropicalis*, Sakayu Shimizu, Koji Yasui, Yoshiki Tani and Hideaki Yamada, Nov. 14, 1979, vol. 91, No. 1, pp. 108-113.

The Regulation of Alanine on the Fermentation of Long-Chain Dicarboxylic Acids in *Candida tropicalis* NPcoN22, Zhou Jianlong, Chiao Juishen, p. 4.

Improved M13 Chage Cloning Vectors and Host Strains; Nucleotide Sequences of the M13mp 18 and pUC19 Vectors, Cleste Yanish-Perron, Jeffrey Viera and Joachim Messing, Gene 33(1985) pp. 103-119.

Inducible Long Chain Alcohol Oxidase from Alkane-grown *Candida tropicalis*, Glenwyn D. Kemp, F. Mark Dickinson and Colin Ratledge, Appl. Microbiol Biotechnol (1988) 29:pp. 370-374.

Aliphatic Hydrocarbons, Matthia Buehler, Joachim Schindler, Chapter 9, pp. 331-384.

[12] One-Step Gene Disruption in Yeast by Rodney J. Rothstein, (1981), pp. 202-211.

Interactions between the .omega.-and .beta.-Oxidations of Fatty Acids, Joseph Vamecoq and Jean-Pierre Draye, Jan. 27, 1987, J. Biochem 102, pp. 225-234.

Production of Macrocyclic Musk Compounds Via Alkanedioic Acids Produced from N-alkanes, L.Hiroshi Okina, Akira Taoka, Namio Uemura, Nov. 20, 1986, pp. 753-760.

Microbial Production of Long-chain Dicarboxylic Acids from n-Alkanes, Part I. Screening and Properties of Microorganism Producing Dicarboxylic Acids, by Isamu Shiio and Ryosuke Uchio, Arg. Biol. Chem. vol. 35, No. 13, pp. 2033-2012 (1971).

Peroxisomal Localization of Enzymes Related to Fatty Acid .beta.-Oxidation in an n-Alkane-grown Yeast, *Candida tropicalis*, Mitsuyoshi Ueda et al., Agric. Biol. Chem. 49(6). pp. 1821-1828 (1985).

Two Acyl-coenzyme A Oxidase in Peroxisomes of the Yeast *Candida tropicalis*: Primary Structures Deduced from Genomic DNA Sequence, Okazaki et al., Proc. Natl. Acad. Sci., USA vol. 83, pp. 1232-1236, Mar. 1986.

Studies of Utilization of Hydrocarbons by Yeasts Part II. Diterminal Oxidation of Alkanes by Yeast, Ogino et al., Agr. Biol. Chem. vol. 29, No. 11, pp. 1009-1015 (1965).

Direct Mutagenesis in *Candida albicans*: On-Step Gene Disruption to Isolate ura3 Mutants, Kelly et al., Molecular and Cellular Biology, Jan. 1987, pp. 199-207.

Studies on the Formation of Long-chain Dicarboxylic Acids from Pure n-alkanes by a Mutant of *Candida tropicalis*, Hill et al., Appl. Microbiol Biotechnol (1986) 24:168-174.

Omega-Hydroxylations, Franz Meussdoerffer pp. 143-146, Biochem. Labs Henkel KGaA.

Degradation of Long-chain n-alkanes by the Yeast *Candida maltosa*, II. Oxidation of n-alkanes and Intermediates Using Microsomal Membrane Fractions, Blasig et al., Appl. Microbiol Biotechnol (1988) 28: pp. 589-597 (1988).

A Method for Gene Disruption that Allows Repeated Use of URA3 Selection in the Construction of Multiply Disrupted Yeast Strains, Alani et al., (1987), Genetics, 116: pp. 541-545.

A Positive Selection for Mutants Lacking Orotidine-5'-phosphate Decarboxylase Activity in Yeast: 5-fluoro-orotic Acid Resistance, Boeckle et al., Mol. Gen. Genet (1984) 197: pp. 345-346.

Mechanisms and Occurrence of Microbial Oxidation of Long-chain Alkanes, Rehm et al. Institute for Microbiologie, pp. 176-217.

Molecular and General Genetics, vol. 214, issued 1988, R. Kelly et al.

"One-step gene disruption by cotransformation to isolate double auxotrophs in *Candida albicans*", pp. 24-31.

Molecular and General Genetics, vol. 217, issued Jan. 1989, M. Kurtz et al,

"Isolation of Hem3 mutants from *Candida albicans* by sequential gene disruption"pp. 47-52.

Molecular and Cell Biology, vol. 7, No. 1, issued Jan. 1987, R. Kelly et al.,

"Directed mutagenesis in *Candida albicans*: One-step gene disruption to isolate ura3 Mutants", pp. 199-207.

Molecular and Cellular Biology, vol. 6, No. 1, issued Jan. 1986, M. Kurtz et al., "Integrative Transformation of *Candida albicans*. Using a Cloned *Candida ADE2* Gene", pp. 142-149.

Proceeding National Academy Science, vol. 83, issued Mar. 1986. K. Okazaki et

al.."Two acyl-coenzyme A oxidases in peroxisomes of the yeast *Candida tropicalis*: Primary structures deduced from genomic DNA sequence pp. 1232-1236.

"Journal of Cell Biology., vol. 105 issued Jul. 1987., T. M. Small et al. "Export of the carboxy-terminal Portion of Acyl-CoA oxidase into Peroxisomes of *Candida tropicalis*" pp. 217-252.

Gene, vol. 58 issued 1987. K. Okazaki et al., "Peroxisomal Acyl-coenzyme A oxidase multigene family of the yeast *Candida tropicalis*: nucleotide sequence of a third gene and its protein product", pp. 37-47.

Journal of Bacteriology, vol. 172, No. 3 issue Aug. 1990, [O. C. Haas et al., "Development of an integrative DNA Transformation System for the yeast *Candida tropicalis*", pp. 4571-4577.

Gene vol. 51, issued 1987, W. W. Murray et al, "The primary structure of a peroxisomal fatty acyl-CoA oxidase from the yeast *Candida tropicalis* p K233", pp. 119-128.

Proceeding National Academy of Science, vol. 76 No. 10 issued Oct. 1979. S. Scherer et al. "Replacement of Chromosome Segments with altered DNA sequences constructed in vitro". pp. 4951-4955.

ART-UNIT: 185

PRIMARY-EXAMINER: Schwartz; Richard A.

ASSISTANT-EXAMINER: LeGuyader; J.

ATTY-AGENT-FIRM: Szoke; Ernest G. Jaeschke; Wayne C. Drach; John E.

ABSTRACT:

The POX genes of *C. tropicalis* are disrupted resulting in the complete blockage of the beta-oxidation pathway in the strain. Fermentation of *C. tropicalis* cells having disrupted genes on alkane, fatty acid and fatty acid ester substrates produces substantially pure dicarboxylic acids in substantially quantitative yield.

30 Claims, 5 Drawing figures

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWIC	Draw. Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	------------	-------

Generate Collection

Terms	Documents
l6 and l4	2

Display

10

Documents, starting with Document:

2

Display Format:

FRO

Change Format

WEST[Generate Collection](#)**Search Results - Record(s) 1 through 2 of 2 returned.**☐ 1. Document ID: US 6020288 A

L9: Entry 1 of 2

File: USPT

Feb 1, 2000

US-PAT-NO: 6020288

DOCUMENT-IDENTIFIER: US 6020288 A

TITLE: Methods and compositions for enhancing cytochrome P450 in plants

DATE-ISSUED: February 1, 2000

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Nonomura; Arthur M.	Boxborough	MA	01719	N/A
Benson; Andrew A.	La Jolla	CA	92037	N/A
Nishio; John N.	Laramie	WY	82070-3917	N/A

US-CL-CURRENT: 504/127; 504/128, 504/130, 504/136, 504/138, 504/140, 504/142,
504/143, 504/144, 504/149

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Draw Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-----------	-------

☐ 2. Document ID: US 5254466 A

L9: Entry 2 of 2

File: USPT

Oct 19, 1993

US-PAT-NO: 5254466

DOCUMENT-IDENTIFIER: US 5254466 A

TITLE: Site-specific modification of the candida tropicals genome

DATE-ISSUED: October 19, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Picataggio; Stephen	Santa Rosa	CA	N/A	N/A
Deanda; Kristine	Graton	CA	N/A	N/A
Eirich; L. Dudley	Santa Rosa	CA	N/A	N/A

US-CL-CURRENT: 435/142; 435/254.22, 435/477, 435/481, 435/483, 435/484,
435/490, 435/6, 435/924

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Draw Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-----------	-------

Generate Collection

Terms	Documents
l8 and l1 and l2 and l6	2

Display 10 Documents, starting with Document: 2

Display Format: CIT Change Format

WEST**Generate Collection****Search Results - Record(s) 1 through 3 of 3 returned.**☐ 1. Document ID: US 6174673 B1

L17: Entry 1 of 3

File: USPT

Jan 16, 2001

US-PAT-NO: 6174673

DOCUMENT-IDENTIFIER: US 6174673 B1

TITLE: High throughput screening for novel enzymes

DATE-ISSUED: January 16, 2001

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Short; Jay M.	Encinitas	CA	N/A	N/A
Keller; Martin	San Diego	CA	N/A	N/A

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Diversa Corporation	San Diego	CA	N/A	N/A	02

APPL-NO: 9/ 098206

DATE FILED: June 16, 1998

PARENT-CASE:

This application is a continuation-in-part of U.S. application Ser. No. 08/876,276, filed Jun. 16, 1997.

INT-CL: [7] C12Q 1/68

US-CL-ISSUED: 435/6; 435/69.1, 435/440, 435/471, 435/476, 435/320.1

US-CL-CURRENT: 435/6; 435/320.1, 435/440, 435/471, 435/476, 435/69.1

FIELD-OF-SEARCH: 435/6, 435/69.1, 435/440, 435/320.1, 435/471, 435/476

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<u>4399219</u>	August 1983	Weaver	435/34
<u>4643968</u>	February 1987	Weaver	435/32
<u>4647536</u>	March 1987	Mosbach et al.	435/177
<u>4916060</u>	April 1990	Weaver	435/29
<u>4959301</u>	September 1990	Weaver et al.	435/5
<u>5055390</u>	October 1991	Weaver et al.	435/5
<u>5225332</u>	July 1993	Weaver et al.	435/29
<u>5824485</u>	October 1998	Thompson et al.	435/6

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
WO 98/56904	December 1998	WOX	
WO 99/49315	September 1999	WOX	
WO99/54494	October 1999	WOX	

OTHER PUBLICATIONS

Zhang et al. FASEB J. vol. 5, pp. 3108-3113, 1991.
Plovins et al. Applied and Environmental Microbiology. vol. 60(12), pp. 4638-4641, 1994.

ART-UNIT: 166

PRIMARY-EXAMINER: Yucel; Remy

ATTY-AGENT-FIRM: Gray Gary Ware & Freidenrich LLP Haile; Lisa A.

ABSTRACT:

Disclosed is a process for identifying clones having a specified activity of interest, which process comprises (i) generating one or more expression libraries derived from nucleic acid directly isolated from the environment; and (ii) screening said libraries utilizing a fluorescence activated cell sorter to identify said clones. More particularly, this is a process for identifying clones having a specified activity of interest by (i) generating one or more expression libraries derived from nucleic acid directly or indirectly isolated from the environment; (ii) exposing said libraries to a particular substrate or substrates of interest; and (iii) screening said exposed libraries utilizing a fluorescence activated cell sorter to identify clones which react with the substrate or substrates. Also provided is a process for identifying clones having a specified activity of interest by (i) generating one or more expression libraries derived from nucleic acid directly or indirectly isolated from the environment; and (ii) screening said exposed libraries utilizing an assay requiring co-encapsulation, a binding event or the covalent modification of a target, and a fluorescence activated cell sorter to identify positive clones.

23 Claims, 18 Drawing figures

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KWC	Draw Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-----------	-------

☐ 2. Document ID: US 6168919 B1

L17: Entry 2 of 3

File: USPT

Jan 2, 2001

US-PAT-NO: 6168919

DOCUMENT-IDENTIFIER: US 6168919 B1

TITLE: Screening methods for enzymes and enzyme kits

DATE-ISSUED: January 2, 2001

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Short; Jay M.	Encinitas	CA	N/A	N/A

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Diversa Corporation	San Diego	CA	N/A	N/A	02

APPL-NO: 8/ 983367

DATE FILED: September 30, 1998

PCT-DATA:

APPL-NO	DATE-FILED	PUB-NO	PUB-DATE	371-DATE	102(E)-DATE
PCT/US96/11854	July 17, 1996	WO97/04077	Feb 6, 1997	Sep 30, 1998	Sep 30, 1998

INT-CL: [7] C12Q 1/68, C12Q 1/00, C12P 19/34, C12N 15/64

US-CL-ISSUED: 435/6; 435/4, 435/91.1, 435/91.4, 435/91.41, 435/252.3, 435/183, 435/320.1, 435/325, 536/23.1, 536/23.2, 536/23.4

US-CL-CURRENT: 435/6; 435/183, 435/252.3, 435/320.1, 435/325, 435/4, 435/91.1, 435/91.4, 435/91.41, 536/23.1, 536/23.2, 536/23.4

FIELD-OF-SEARCH: 435/4, 435/6, 435/91.1, 435/91.4, 435/91.41, 435/252.3, 435/183, 435/320.1, 435/325, 536/23.1, 536/23.2, 536/23.4

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<u>5171684</u>	December 1992	Yen et al.	435/252.3
<u>5712146</u>	January 1998	Khosla et al.	435/252.35
<u>5958672</u>	September 1999	Short	435/4

OTHER PUBLICATIONS

Lactic Dehydrogenase, Sigma catalog, p. 634, 1997.

Anderson et al., Met. Enzymol., vol. 68, pp. 428-436, 1979.

Promega Catalog, p. 205, 1993.

Sambrook et al., Molecular Cloning, Cold Spring Harbor Laboratory Press, vol. 2, p. 9.30 and pp. 12.1-12.20, 1989.*

ART-UNIT: 162

PRIMARY-EXAMINER: Achutamurthy; Ponnathapu

ASSISTANT-EXAMINER: Tung; Peter P.

ATTY-AGENT-FIRM: Gray, Cary, Ware & Freidenrich LLP Haile; Lisa A.

ABSTRACT:

Recombinant enzyme libraries and kits where a plurality of enzymes are each characterized by different physical and/or chemical characteristics and classified by common characteristics. The characteristics are determined by screening of recombinant enzymes expressed by a DNA library produced from various microorganisms. Also disclosed is a process for identifying clones of a recombinant library which express a protein with a desired ctivity by screening a library of expression clones randomly produced from DNA of at least one microorganism, said screeing being effected on expression products of said clones to thereby identify clones which express a protein with a desired activity. Also disclosed is a process of screening clones having DNA from an uncultivated microorganism for a specified protein activity by screening for a specified protein activity in a library of clones prepared by (i) recovering DNA from a DNA population derived from at least one uncultivated microorganism; and (ii) transforming a host with recovered DNA to produce a library of clones which is screened for the specified protein activity.

9 Claims, 8 Drawing figures

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMC	Draw Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	-----	-----------	-------

☐ 3. Document ID: US 5254466 A

L17: Entry 3 of 3

File: USPT

Oct 19, 1993

US-PAT-NO: 5254466

DOCUMENT-IDENTIFIER: US 5254466 A

TITLE: Site-specific modification of the candida tropicals genome

DATE-ISSUED: October 19, 1993

INVENTOR-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY
Picataggio; Stephen	Santa Rosa	CA	N/A	N/A
Deanda; Kristine	Graton	CA	N/A	N/A
Eirich; L. Dudley	Santa Rosa	CA	N/A	N/A

ASSIGNEE-INFORMATION:

NAME	CITY	STATE	ZIP CODE	COUNTRY	TYPE CODE
Henkel Research Corporation	Santa Rosa	CA	N/A	N/A	02

APPL-NO: 7/ 432091

DATE FILED: November 6, 1989

INT-CL: [5] C12N 15/09, C12P 7/44

US-CL-ISSUED: 435/142; 435/172.1, 435/172.3, 435/924, 435/254.22

US-CL-CURRENT: 435/142; 435/254.22, 435/477, 435/481, 435/483, 435/484,
435/490, 435/6, 435/924

FIELD-OF-SEARCH: 435/67.1, 435/142, 435/172.3, 435/172.1, 435/940, 435/255,
435/924, 935/22, 935/28

PRIOR-ART-DISCLOSED:

U.S. PATENT DOCUMENTS

PAT-NO	ISSUE-DATE	PATENTEE-NAME	US-CL
<u>4735901</u>	April 1988	Kurtz et al.	435/172.3

FOREIGN PATENT DOCUMENTS

FOREIGN-PAT-NO	PUBN-DATE	COUNTRY	US-CL
183070	April 1986	EPX	

OTHER PUBLICATIONS

Okazaki et al. Gene 57:37-44 (1987).
Okazaki et al. Biochem. 83:1232-1236 (1986).
Kelly et al. Mol. Cell. Biol. Jan 1987: p. 199-207.
Gillam et al. Mol. Gen. Genet. 198: 179 (1984).
Stromnaes et al. J. Bact. Jan. 1986: 197.
Ditchburn et al. J. Gen Microbiol. 67:299 (1971).

Gaillasdin et al. Chem Abst. vol. 77 (23) No. 149584R 1971.
Watanabe et al. Chem. Abst. vol. 70 (15) No. 65372n 1968.
Small et al. J. Cell. Biol. 105:247 (1987).
Strain and Species Identification by Restriction . . . Candida Species, Journal of Bacteriology, Apr. 1987, pp. 1639-1643, Magee et al.
The Genetics of Candida, CRC Press, 1990, pp. 177-186, Kirsch et al.
Genetics of Candida albicans, Microbiological Reviews, Sep. 1990, pp. 226-241, Scherer et al.
Isolation and Determination of Yeasts . . . Source of Carbon, Agr. Biol. Chem., vol. 30, No. 12, pp. 1175-1182, 1966, Tanabe et al.
Methods for the Genetics and Molecular Biology of Candida albicans, Analytical Biochemistry, 175, pp. 361-372 (1988), Magee et al.
The Genus Candida Berkhout nom. conserv. . . . Delimitation, System. Appl. Microbiol. 12, pp. 183-190 (1989), Viljoen et al.
Variation in the electrophoretic karyotype . . . in Candida albicans, Journal of General Microbiology (1990), 136, pp. 2433-2442, Iwaguchi et al.
The Carboxyl-terminal Tripeptide Ala-Lys-Ile . . . Yeast Peroxisomes, The Journal of Biological Chemistry, vol. 266, No. 34, pp. 23197-23208, 1991, Aitchison et al.
In vivo import of Candida tropicalis . . . Candida albicans, Current Genetics, 1990, 17:481-486, Aitchison et al.
Redefinition of Candida Berkhout and the consequent emendation of Cryptococcus Kuetzing and Rhodotorula Harrison, accepted 31 Mar. 1988, Weijman et al.
Interspecific Complementation Analysis . . . Candida albicans Adenine Auxotrophs, Journal of Bacteriology, Jun. 1989, vol. 171, No. 6, pp. 3586-3589, Corner et al.
Sequence and transcript analysis . . . -phosphate decarboxylase, Curr Genet (1989), 16:153-157, Losberger et al.
The yeasts a taxonomic study, third edition, Elsevier Science Publishers B.V.--Amsterdam, N.J.W. Kreger-van Rij, ed.
Differential Identification of Candida Species . . . Polypeptide Profiles, Analytical Biochemistry 175, 548-551 (1988), Shen et al.
Transformation of Intact Yeast Cells Treated with Alkali Cations, Hisao Ito, Yasuki Fukuda, Kousaku Murata and Akira Kimura, Jan. 1983, pp. 163-168.
Acyl-CoA Oxidase From Candida tropicalis, Sakayu Shimizu, Koji Yasui, Yoshiki Tani and Hideaki Yamada, Nov. 14, 1979, vol. 91, No. 1, pp. 108-113.
The Regulation of Alanine on the Fermentation of Long-Chain Dicarboxylic Acids in Candida tropicalis NPcoN22, Zhou Jianlong, Chiao Juishen, p. 4.
Improved M13 Chage Cloning Vectors and Host Strains; Nucleotide Sequences of the M13mp 18 and pUC19 Vectors, Cleste Yanish-Perron, Jeffrey Viera and Joachim Messing, Gene 33(1985) pp. 103-119.
Inducible Long Chain Alcohol Oxidase from Alkane-grown Candida tropicalis, Glenwyn D. Kemp, F. Mark Dickinson and Colin Ratledge, Appl. Microbiol Biotechnol (1988) 29:pp. 370-374.
Aliphatic Hydrocarbons, Matthia Buehler, Joachim Schindler, Chapter 9, pp. 331-384.
[12] One-Step Gene Disruption in Yeast by Rodney J. Rothstein, (1981), pp. 202-211.
Interactions between the .omega.-and .beta.-Oxidations of Fatty Acids, Joseph Vamecoq and Jean-Pierre Draye, Jan. 27, 1987, J. Biochem 102, pp. 225-234.
Production of Macrocyclic Musk Compounds Via Alkanedioic Acids Produced from N-alkanes, LHiroshi Okina, Akira Taoka, Namio Uemura, Nov. 20, 1986, pp. 753-760.
Mircobial Production of Long-chain Dicarboxylic Acids from n-Alkanes, Part I. Screening and Properties of Microorganism Producing Dicarboxylic Acids, by Isamu Shiio and Ryosuke Uchio, Arg. Biol. Chem. vol. 35, No. 13, pp. 2033-2012 (1971).
Peroxidomal Localization of Enzymes Related to Fatty Acid .beta.-Oxidation in an n-Alkane-grown Yeast, Candida tropicalis, Mitsuyoshi Ueda et al., Agric. Biol. Chem. 49(6). pp. 1821-1828 (1985).
Two Acyl-coenzyme A Oxidase in Peroxisomes of the Yeast Candida tropicalis: Primary Structures Deduced from Genomic DNA Sequence, Okazaki et al., Proc. Natl. Acad. Sci., USA vol. 83, pp. 1232-1236, Mar. 1986.
Studies of Utilization of Hydrocarbons by Yeasts Part II. Diterminal Oxidation of Alkanes by Yeast, Ogino et al., Agr. Biol. Chem. vol. 29, No. 11, pp.

1009-1015 (1965).

Direct Mutagenesis in *Candida albicans*: On-Step Gene Disruption to Isolate *ura3* Mutants, Kelly et al., *Molecular and Cellular Biology*, Jan. 1987, pp. 199-207. Studies on the Formation of Long-chain Dicarboxylic Acids from Pure n-alkanes by a Mutant of *Candida tropicalis*, Hill et al., *Appl. Microbiol Biotechnol* (1986) 24:168-174.

Omega-Hydroxylations, Franz Meussdoerffer pp. 143-146, *Biochem. Labs Henkel KGaA*.

Degradation of Long-chain n-alkanes by the Yeast *Candida maltosa*, II. Oxidation of n-alkanes and Intermediates Using Microsomal Membrane Fractions, Blasig et al., *Appl. Microbiol Biotechnol* (1988) 28: pp. 589-597 (1988).

A Method for Gene Disruption that Allows Repeated Use of *URA3* Selection in the Construction of Multiply Disrupted Yeast Strains, Alani et al., (1987), *Genetics*, 116: pp. 541-545.

A Positive Selection for Mutants Lacking Orotidine-5'-phosphate Decarboxylase Activity in Yeast: 5-fluoro-orotic Acid Resistance, Boeckle et al., *Mol. Gen. Genet* (1984) 197: pp. 345-346.

Mechanisms and Occurrence of Microbial Oxidation of Long-chain Alkanes, Rehm et al. *Institute for Microbiologie*, pp. 176-217.

Molecular and General Genetics, vol. 214, issued 1988, R. Kelly et al.

"One-step gene disruption by cotransformation to isolate double auxotrophs in *Candida albicans*", pp. 24-31.

Molecular and General Genetics, vol. 217, issued Jan. 1989, M. Kurtz et al.,

"Isolation of *Hem3* mutants from *Candida albicans* by sequential gene disruption" pp. 47-52.

Molecular and Cell Biology, vol. 7, No. 1, issued Jan. 1987, R. Kelly et al.,

"Directed mutagenesis in *Candida albicans*: One-step gene disruption to isolate *ura3* Mutants", pp. 199-207.

Molecular and Cellular Biology, vol. 6, No. 1, issued Jan. 1986, M. Kurtz et al., "Integrative Transformation of *Candida albicans*. Using a Cloned *Candida ADE2* Gene", pp. 142-149.

Proceeding National Academy Science, vol. 83, issued Mar. 1986. K. Okazaki et al. "Two acyl-coenzyme A oxidases in peroxisomes of the yeast *Candida tropicalis*: Primary structures deduced from genomic DNA sequence pp. 1232-1236.

"*Journal of Cell Biology*., vol. 105 issued Jul. 1987., T. M. Small et al.

"Export of the carboxy-terminal Portion of Acyl-CoA oxidase into Peroxisomes of *Candida tropicalis*" pp. 217-252.

Gene, vol. 58 issued 1987. K. Okazaki et al., "Peroxisomal Acyl-coenzyme A oxidase multigene family of the yeast *Candida tropicalis*: nucleotide sequence of a third gene and its protein product", pp. 37-47.

Journal of Bacteriology, vol. 172, No. 3 issue Aug. 1990, [O. C. Haas et al., "Development of an integrative DNA Transformation System for the yeast *Candida tropicalis*", pp. 4571-4577.

Gene vol. 51, issued 1987, W. W. Murray et al, "The primary structure of a peroxisomal fatty acyl-CoA oxidase from the yeast *Candida tropicalis* p K233", pp. 119-128.

Proceeding National Academy of Science, vol. 76 No. 10 issued Oct. 1979. S. Scherer et al. "Replacement of Chromosome Segments with altered DNA sequences constructed in vitro". pp. 4951-4955.

ART-UNIT: 185

PRIMARY-EXAMINER: Schwartz; Richard A.

ASSISTANT-EXAMINER: LeGuyader; J.

ATTY-AGENT-FIRM: Szoke; Ernest G. Jaeschke; Wayne C. Drach; John E.

ABSTRACT:

The POX genes of *C. tropicalis* are disrupted resulting in the complete blockage of the beta-oxidation pathway in the strain. Fermentation of *C. tropicalis* cells having disrupted genes on alkane, fatty acid and fatty acid ester substrates produces substantially pure dicarboxylic acids in substantially quantitative yield.

30 Claims, 5 Drawing figures

Full	Title	Citation	Front	Review	Classification	Date	Reference	Claims	KMIC	Draw. Desc	Image
------	-------	----------	-------	--------	----------------	------	-----------	--------	------	------------	-------

Generate Collection

Terms	Documents
(116 or 115 or 114 or 113 or 112 or 111 or 110) and 18 and 16 and 11	3

Display 10 Documents, starting with Document: 3

Display Format: FRO Change Format

=> d his full

(FILE 'HOME' ENTERED AT 08:16:29 ON 10 MAY 2001)

FILE 'CAPLUS' ENTERED AT 08:16:37 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 08:16:41 ON 10 MAY 2001

L1 1 SEA ABB=ON PLU=ON 62213-32-5/RN
 D
L2 1 SEA ABB=ON PLU=ON 9038-14-6/RN
 D
L3 1 SEA ABB=ON PLU=ON 9023-03-4/RN
 D

FILE 'CAPLUS' ENTERED AT 08:18:49 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 08:19:02 ON 10 MAY 2001

 SET SMARTSELECT ON
L4 SEL PLU=ON L2 1- CHEM : 26 TERMS
 SET SMARTSELECT OFF

FILE 'CAPLUS' ENTERED AT 08:19:03 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 08:19:03 ON 10 MAY 2001

 SET SMARTSELECT ON
L5 SEL PLU=ON L3 1- CHEM : 16 TERMS
 SET SMARTSELECT OFF

FILE 'CAPLUS' ENTERED AT 08:19:04 ON 10 MAY 2001

L6 15271 SEA ABB=ON PLU=ON L4
L7 7478 SEA ABB=ON PLU=ON L5
L8 1165 SEA ABB=ON PLU=ON L6 AND L7
L9 188643 SEA ABB=ON PLU=ON DICARBOXYLIC ACID# OR (CARBOXYLIC ACIDS
 (L) DICARBOXYLIC) OR CARBOXYLIC ACID# OR MONOCARBOXYLIC ACID#
L10 0 SEA ABB=ON PLU=ON L8 (L) L9
L11 5883 SEA ABB=ON PLU=ON (HYDROXYLATION (L) .OMEGA.-HYDROXYLATION)
 OR (HYDROXYLATION (L) ALKANE#) OR (.ALPHA.-HYDROXYLATION) OR
 (HYDROXYLATION (L) .ALPHA.)
L12 0 SEA ABB=ON PLU=ON L9 AND L10
L13 149 SEA ABB=ON PLU=ON L9 AND L11
L14 8 SEA ABB=ON PLU=ON L13 AND (L6 OR L7)
L15 8 DUP REM L14 (0 DUPLICATES REMOVED)
 D IBIB AB 1-8
 D IBIB AB HIT 1-8
L16 8 SEA L15
L17 5 SEA ABB=ON PLU=ON L16 AND (PREP/RL OR PREPAR? OR SYNTH? OR
 MAK?)
 D IBIB AB HIT 1-5

FILE HOME

FILE CAPLUS

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications.

FILE COVERS 1947 - 10 May 2001 VOL 134 ISS 20

FILE LAST UPDATED: 9 May 2001 (20010509/ED)

This file contains CAS Registry Numbers for easy and accurate substance identification.

This file supports REGISTRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

CAPLUS now provides online access to patents and literature covered in CA from 1947 to the present. On April 22, 2001, bibliographic information and abstracts were added for over 2.2 million references published in CA from 1947 to 1966.

The CA Lexicon is now available in the Controlled Term (/CT) field. Enter HELP LEXICON for full details.

Attention, the CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE REGISTRY

STRUCTURE FILE UPDATES: 9 MAY 2001 HIGHEST RN 335078-44-9
DICTIONARY FILE UPDATES: 9 MAY 2001 HIGHEST RN 335078-44-9

TSCA INFORMATION NOW CURRENT THROUGH January 11, 2001.

Please note that search-term pricing does apply when conducting SmartSELECT searches.

Structure search limits have been increased. See HELP SLIMIT for details.

=> d ibib ab hit 1-5

L17 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 2000:133824 CAPLUS

DOCUMENT NUMBER: 132:162018

TITLE: DNA shuffling of **monooxygenase** genes for
production of industrial chemicals

INVENTOR(S): Affholter, Joseph A.; Davis, Christopher; Selifonov,
Sergey A.

PATENT ASSIGNEE(S): Maxygen, Inc., USA

SOURCE: PCT Int. Appl., 153 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000009682	A1	20000224	WO 1999-US18424	19990812
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
AU 9953479	A1	20000306	AU 1999-53479	19990812
PRIORITY APPLN. INFO.:			US 1998-96271	P 19980812
			US 1999-130810	P 19990423
			WO 1999-US18424	W 19990812

OTHER SOURCE(S): MARPAT 132:162018

AB This invention provides improved **monooxygenases**, dehydrogenases, and transferases that are useful for the biocatalytic **synthesis** of compds. such as .alpha.-hydroxycarboxylic acids, and aryl- and alkyl-, hydroxy compds. The polypeptides provided herein are improved in properties such as regioselectivity, enzymic activity, stereospecificity, and the like. Methods for obtaining recombinant polynucleotides that encode these improved polypeptides are also provided, as are organisms that express the polypeptides and are thus useful for carrying out said biocatalytic **syntheses**. In the methods for obtaining **monooxygenase** genes, a plurality of parental forms (homologs) of a selected nucleic acid are recombined. The selected nucleic acid derived either from one or more parental nucleic acid(s) which encodes a **monooxygenase** enzyme, or a fragment thereof, or from a parental nucleic acid which does not encode **monooxygenase**, but which is a candidate for DNA shuffling to develop **monooxygenase** activity. The plurality of forms of the selected nucleic acid differ from each

other

in at lease one (and typically two or more) nucleotides, and, upon recombination, provide a library of recombinant **monooxygenase** nucleic acids. The library can be an in vitro set of mols., or present

in

cells, phage or the like. The library is screened to identity at least one recombinant **monooxygenase** nucleic acid that exhibits distinct or improved **monooxygenase** activity compared to the parental nucleic acid or nucleic acids. Also provided by the invention

are methods for increasing said solvent resistance of organisms that are used in the **synthetic** methods.

REFERENCE COUNT: 18
REFERENCE(S): (1) Affymax Tech Nv; WO 9720078 A 1997 CAPLUS
(2) Agency Of Ind Sci & Technology; JP 05-049474 A 1993 CAPLUS
(3) Aoyama, T; JOURNAL OF BIOLOGICAL CHEMISTRY 1989, V264(18), P10388 CAPLUS
(4) Cramer, A; NATURE 1998, V391, P288 CAPLUS
(5) Dierks, E; THE JOURNAL OF BIOLOGICAL CHEMISTRY 1998, V273(36), P23055 CAPLUS

ALL CITATIONS AVAILABLE IN THE RE FORMAT

TI DNA shuffling of **monooxygenase** genes for production of industrial chemicals

AB This invention provides improved **monooxygenases**, dehydrogenases, and transferases that are useful for the biocatalytic **synthesis** of compds. such as .alpha.-hydroxycarboxylic acids, and aryl- and alkyl-, hydroxy compds. The polypeptides provided herein are improved in properties such as regioselectivity, enzymic activity, stereospecificity, and the like. Methods for obtaining recombinant polynucleotides that encode these improved polypeptides are also provided, as are organisms that express the polypeptides and are thus useful for carrying out said biocatalytic **syntheses**. In the methods for obtaining **monooxygenase** genes, a plurality of parental forms (homologs) of a selected nucleic acid are recombined. The selected nucleic acid derived either from one or more parental nucleic acid(s) which encodes a **monooxygenase** enzyme, or a fragment thereof, or from a parental nucleic acid which does not encode **monooxygenase**, but which is a candidate for DNA shuffling to develop **monooxygenase** activity. The plurality of forms of the selected nucleic acid differ from each other in at least one (and typically two or more) nucleotides, and, upon recombination, provide a library of recombinant **monooxygenase** nucleic acids. The library can be an in vitro set of mols., or present in cells, phage or the like. The library is screened to identify at least one recombinant **monooxygenase** nucleic acid that exhibits distinct or improved **monooxygenase** activity compared to the parental nucleic acid or nucleic acids. Also provided by the invention are methods for increasing said solvent resistance of organisms that are used in the **synthetic** methods.

ST **monooxygenase** industrial **synthesis** DNA shuffling gene

IT Bioreactors
Nucleic acid library
Phage display library
Recombination, genetic
Regiochemistry
(DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Gene
RL: BPR (Biological process); BUU (Biological use, unclassified); BIOL (Biological study); PROC (Process); USES (Uses)
(DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Amines, biological studies
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(N-dealkylation; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Ethers, biological studies

RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(O-dealkylation; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Thiols (organic), biological studies
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(S-dealkylation; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Ethers, biological studies
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(allyl, substrate; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Baeyer-Villiger oxidation
Dealkylation
Decarboxylation
Dehalogenation
Dehydrogenation
Epoxidation
Hydroxylation
Oxidation
(enzymic; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Alkenes, biological studies
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(epoxidn.; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Aromatic hydrocarbons, biological studies
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(halo, oxidative dehalogenation; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT **Carboxylic acids**, biological studies
RL: BPN (Biosynthetic preparation); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); **PREP (Preparation)**
(hydroxy, conversion from olefins; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT **Alkanes**, biological studies
Aromatic hydrocarbons, biological studies
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(**hydroxylation**; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Chemicals
(industrial; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Solvents
(org., resistance to; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Sulfonylureas
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
(oxygenation; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)

IT Proteins, specific or class
RL: BPN (Biosynthetic preparation); BIOL (Biological study); **PREP (Preparation)**

- (solvent-resistant; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT Glycols, biological studies
 RL: BPN (Biosynthetic preparation); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); **PREP (Preparation)**
 (vicinal, conversion from olefins; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 9003-99-0P, Peroxidase 9013-18-7P, Acyl-CoA Ligase 9023-26-1P, Coenzyme A transferase 9024-04-8P, Mandelate racemase 9031-56-5P, Ligase 9033-07-2P, Glycosyltransferase 9033-25-4P, Methyltransferase 9035-51-2P, Cytochrome P 450, **preparation** 9035-82-9P, Dehydrogenase **9038-14-6P, Monooxygenase** 9047-61-4P, Transferase 9048-63-9P, Epoxide hydrolase 9054-54-0P, Acyltransferase 9080-22-2P, Racemase 102055-73-2P, Oxo acid decarboxylase
 RL: BPN (Biosynthetic preparation); CAT (Catalyst use); BIOL (Biological study); **PREP (Preparation)**; USES (Uses)
 (DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 60-12-8DP, 2-Phenylethanol, substituted 60-12-8P, 2-Phenylethanol 100-51-6DP, Benzyl alcohol, substituted 100-51-6P, Benzyl alcohol, biological studies 122-97-4DP, 3-Phenylpropanol, substituted 122-97-4P, 3-Phenylpropanol
 RL: BPN (Biosynthetic preparation); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); **PREP (Preparation)**
 (DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 145-13-1P, Pregnenolone
 RL: BPN (Biosynthetic preparation); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); **PREP (Preparation)**
 (conversion from cholesterol; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 81093-37-0P, Pravastatin
 RL: BPN (Biosynthetic preparation); MFM (Metabolic formation); BIOL (Biological study); FORM (Formation, nonpreparative); **PREP (Preparation)**
 (conversion from mevastatin; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 57-88-5, Cholesterol, biological studies
 RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
 (conversion to pregnenolone; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 119-84-6, 3,4-Dihydrocoumarin 73573-88-3, Mevastatin
 RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
 (hydroxylation; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 59865-13-3, Cyclosporin 79217-60-0, Cyclosporin
 RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study); PROC (Process)
 (modification; DNA shuffling of **monooxygenase** genes for prodn. of industrial chems.)
- IT 78-79-5, Isoprene, biological studies 100-42-5, Styrene, biological studies 106-99-0, Butadiene, biological studies 107-02-8, Acrolein, biological studies 557-40-4, Diallyl ether 1321-74-0, Divinylbenzene, biological studies 1321-74-0D, Divinylbenzene, substituted 1746-13-0, Allyl phenyl ether 1746-13-0D, Allyl phenyl ether, substituted

25168-07-4, Vinylcyclohexene 40356-67-0, Vinylnorbornene
RL: BPR (Biological process); RCT (Reactant); BIOL (Biological study);
PROC (Process)
(substrate; DNA shuffling of **monooxygenase** genes for prodn.
of industrial chems.)

L17 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1999:77701 CAPLUS

DOCUMENT NUMBER: 130:138397

TITLE: Yeasts with elevated cytochrome P450 levels and their
use in the manufacture of monoterminial and diterminial
aliphatic carboxylates from alkanes

INVENTOR(S): Fallon, Robert D.; Payne, Mark S.; Picataggio,
Stephen

PATENT ASSIGNEE(S): K.; Wu; Shijun
E.I. Du Pont De Nemours and Company, USA

SOURCE: PCT Int. Appl., 58 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9904014	A2	19990128	WO 1998-US14935	19980720
WO 9904014	A3	19990520		
W: AU, CA, IL, JP, NZ, US				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9884982	A1	19990210	AU 1998-84982	19980720
EP 1003881	A2	20000531	EP 1998-935806	19980720
R: DE, DK, ES, FR, GB, IT, NL, SE, IE				
PRIORITY APPLN. INFO.:		US 1997-53215	P	19970721
		WO 1998-US14935	W	19980720
AB	Strains of Pichia pastoris and Candida maltosa that have increased cytochrome P 450 activity or lack the .beta.-oxidn. pathway that can be used to convert c6-C22 alkanes to monoterminial and diterminial carboxylic acids are described. Expression constructs carrying cytochrome P 450 genes under the control of powerful inducible promoters (AOX1, PGK) were constructed by std. methods and introduced into P. pastoris or C. maltosa to give novel or increased P 450 activities. Disruption of .beta.-oxidn. in C. maltosa was achieved by insertional inactivation of the POX4 gene. A strain of C. maltosa cultured in a complete medium that was supplemented with dodecane 20 g/L when glucose was almost completely depleted was able to generate dodecanedioic acid at 3.4 g/h for 51 h.			
AB	Strains of Pichia pastoris and Candida maltosa that have increased cytochrome P 450 activity or lack the .beta.-oxidn. pathway that can be used to convert c6-C22 alkanes to monoterminial and diterminial carboxylic acids are described. Expression constructs carrying cytochrome P 450 genes under the control of powerful inducible promoters (AOX1, PGK) were constructed by std. methods and introduced into P. pastoris or C. maltosa to give novel or increased P 450 activities. Disruption of .beta.-oxidn. in C. maltosa was achieved by insertional inactivation of the POX4 gene. A strain of C. maltosa cultured in a complete medium that was supplemented with dodecane 20 g/L when glucose was almost completely depleted was able to generate dodecanedioic acid at			

3.4 g/h for 51 h.

ST Pichia **alkane hydroxylation carboxylic dicarboxylic acid** manuf; cytochrome P450 Candida Pichia **alkane carboxylic acid** fermn; Candida **alkane hydroxylation carboxylic dicarboxylic acid** manuf

IT **Carboxylic acids, preparation**
Dicarboxylic acids
 RL: BMF (Bioindustrial manufacture); BIOL (Biological study); **PREP**
(Preparation)
 (C6-22; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

IT Fermentation
 (**carboxylic acids**, with transgenic yeasts; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

IT Plasmid vectors
 (pLPA1T, alkane **monooxygenase** gene on, expression in Pichia of; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

IT Plasmid vectors
 (pSW84, alkane **monooxygenase** and cytochrome reductase genes on, expression in Candida of; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

IT Plasmid vectors
 (pSW87, alkane **monooxygenase** and cytochrome reductase genes on, expression in Candida of; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

IT **9038-14-6, Monooxygenase** 9059-16-9, Alkane **monooxygenase** 106178-16-9, Fatty acid **monooxygenase**
 RL: BPR (Biological process); BUU (Biological use, unclassified); CAT (Catalyst use); BIOL (Biological study); PROC (Process); USES (Uses)
 (in alkane utilization by yeasts; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

IT 693-23-2P, Dodecanedioic acid
 RL: BMF (Bioindustrial manufacture); BIOL (Biological study); **PREP**
(Preparation)
 (manuf. from dodecane of; yeasts with elevated cytochrome P 450 levels and their use in manuf. of monoterminial and diterminial aliph. carboxylates from alkanes)

L17 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1993:5685 CAPLUS

DOCUMENT NUMBER: 118:5685

TITLE: Metabolic engineering of Candida tropicalis for the production of long-chain **dicarboxylic acids**

AUTHOR(S): Picataggio, Stephen; Rohrer, Tracy; Deanda, Kristine; Lanning, Dawn; Reynolds, Robert; Mielenz, Jonathan; Eirich, L. Dudley

CORPORATE SOURCE: Microb. Technol. Dep., Cognis Inc., Santa Rosa, CA, 95407, USA

SOURCE: Bio/Technology (1992), 10(8), 894-8
 CODEN: BTCHDA; ISSN: 0733-222X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An industrial strain of the yeast *C. tropicalis* was engineered for the efficient prodn. of long-chain **dicarboxylic acids**, which are important raw materials for the chem. industry. By sequential disruption of the 4 genes encoding both isoenzymes of the acyl-CoA oxidase

which catalyzes the first reaction in the .beta.-oxidn. pathway, **alkane** and fatty acid substrates were successfully redirected to the .omega.-oxidn. pathway. Consequently, the conversion efficiency and chem. selectivity of their terminal oxidn. to the corresponding **dicarboxylic acids** was improved to 100%. The specific productivity of the bioconversion was increased further by amplification of the **cytochrome P 450**

monooxygenase and NADPH-cytochrome reductase genes encoding the rate-limiting .omega.-hydroxylase in the .omega.-oxidn. pathway. The amplified strains demonstrated increased .omega.-hydroxylase activity and a 30% increase in productivity compared to the .beta.-oxidn.-blocked strain in fermns. The bioconversion is effective for the selective terminal oxidn. of both satd. and unsatd. linear aliph. substrates with C12-22 chain-lengths and also avoids the undesirable chain modifications assocd. with passage through the .beta.-oxidn. pathway, such as unsatn., **hydroxylation**, or chain shortening. It is now possible to efficiently produce a wide range of previously unavailable satd. and unsatd. **dicarboxylic acids** with a high degree of purity.

TI Metabolic engineering of *Candida tropicalis* for the production of long-chain **dicarboxylic acids**

AB An industrial strain of the yeast *C. tropicalis* was engineered for the efficient prodn. of long-chain **dicarboxylic acids**, which are important raw materials for the chem. industry. By sequential disruption of the 4 genes encoding both isoenzymes of the acyl-CoA oxidase

which catalyzes the first reaction in the .beta.-oxidn. pathway, **alkane** and fatty acid substrates were successfully redirected to the .omega.-oxidn. pathway! Consequently, the conversion efficiency and chem. selectivity of their terminal oxidn. to the corresponding **dicarboxylic acids** was improved to 100%. The specific productivity of the bioconversion was increased further by amplification of the **cytochrome P 450**

monooxygenase and NADPH-cytochrome reductase genes encoding the rate-limiting .omega.-hydroxylase in the .omega.-oxidn. pathway. The amplified strains demonstrated increased .omega.-hydroxylase activity and a 30% increase in productivity compared to the .beta.-oxidn.-blocked strain in fermns. The bioconversion is effective for the selective terminal oxidn. of both satd. and unsatd. linear aliph. substrates with C12-22 chain-lengths and also avoids the undesirable chain modifications assocd. with passage through the .beta.-oxidn. pathway, such as unsatn., **hydroxylation**, or chain shortening. It is now possible to efficiently produce a wide range of previously unavailable satd. and unsatd. **dicarboxylic acids** with a high degree of purity.

ST genetic engineering *Candida* **dicarboxylic acid** prodn

IT *Candida tropicalis*
(genetic engineering of, for long-chain **dicarboxylic acid** prodn.)

IT Genetic engineering
(of *Candida tropicalis*, for long-chain **dicarboxylic acid** prodn.)

IT **Carboxylic acids, preparation**

RL: PREP (Preparation)

(di-, long-chain, manuf. of, genetic engineering of *Candida tropicalis* for)

L17 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1985:61628 CAPLUS

DOCUMENT NUMBER: 102:61628

TITLE: Oxidative ether cleavage with p-nitroperbenzoic acid

AUTHOR(S): Schneider, Hans Joerg; Ahlhelm, Alfred; Mueller, Walter

CORPORATE SOURCE: Univ. Saarlandes, Saarbruecken, D-6600/11, Fed. Rep. Ger.

SOURCE: Chem. Ber. (1984), 117(11), 3297-302

CODEN: CHBEAM; ISSN: 0009-2940

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The reaction of p-nitroperbenzoic acid in CHCl_3 with $\text{Me}_2\text{CH}(\text{CH}_2)_3\text{OMe}$ and $(\text{Me}_2\text{CHCH}_2\text{CH}_2)_2\text{O}$ leads by selective attack at C-H bonds in the .

alpha.-position to the ether oxygen to hemiacetals, which decomp.

to aldehydes and alcs., yielding **carboxylic acids**.

Secondary alkoxy groups, as in ethoxycyclohexane, furnish Baeyer-Villiger oxidn. products of initially formed ketones. Kinetic measurements with substituted benzyl Me ethers show a Hammett reaction const. $\rho = -0.9$, which is in accord with the obsd. relatively small discrimination between secondary and tertiary C-H bonds. The results are compared with similar **hydroxylations** of **alkanes** and with **monooxygenase** reactions and point to oxenoid transition states. Radical reactions, as found with some **alkanes** are not obsd., which is shown by the small amts. of PhNO_2 (1.0%) formed during the reaction. ^{13}C -NMR shifts of several ethers and oxidn. products are reported.

AB The reaction of p-nitroperbenzoic acid in CHCl_3 with $\text{Me}_2\text{CH}(\text{CH}_2)_3\text{OMe}$ and $(\text{Me}_2\text{CHCH}_2\text{CH}_2)_2\text{O}$ leads by selective attack at C-H bonds in the .

alpha.-position to the ether oxygen to hemiacetals, which decomp.

to aldehydes and alcs., yielding **carboxylic acids**.

Secondary alkoxy groups, as in ethoxycyclohexane, furnish Baeyer-Villiger oxidn. products of initially formed ketones. Kinetic measurements with substituted benzyl Me ethers show a Hammett reaction const. $\rho = -0.9$, which is in accord with the obsd. relatively small discrimination between secondary and tertiary C-H bonds. The results are compared with similar **hydroxylations** of **alkanes** and with **monooxygenase** reactions and point to oxenoid transition states. Radical reactions, as found with some **alkanes** are not obsd., which is shown by the small amts. of PhNO_2 (1.0%) formed during the reaction. ^{13}C -NMR shifts of several ethers and oxidn. products are reported.

IT 123-51-3P 502-44-3P 503-74-2P 622-45-7P 626-89-1P 646-07-1P

2412-73-9P 5299-60-5P 94368-15-7P 94368-16-8P

RL: PREP (Preparation)

(formation and carbon-13 NMR of)

L17 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1977:498893 CAPLUS

DOCUMENT NUMBER: 87:98893

TITLE: Biosynthesis of cutin. **omega.**-

Hydroxylation of fatty acids by a microsomal **preparation** from germinating *Vicia faba*

AUTHOR(S): Soliday, Charles L.; Kolattukudy, P. E.

CORPORATE SOURCE: Dep. Agric. Chem., Washington State Univ., Pullman, Wash., USA

SOURCE: Plant Physiol. (1977), 59(6), 1116-21

CODEN: PLPHAY

DOCUMENT TYPE: Journal
LANGUAGE: English

AB **.omega.-Hydroxylation** of fatty acids, which is a key reaction in the biosynthesis of cutin and suberin, was demonstrated in a cell-free **prepn.** from a higher plant. A crude microsomal fraction (105,000-g pellet) from germinating embryonic shoots of *V. faba* catalyzed the conversion of palmitic acid to **.omega.-hydroxypalmitic**

acid.

As the crude cell-free **prepn.** also catalyzes the formation of other hydroxy acids such as **.alpha.-** and **.beta.-hydroxy** acids, the **.omega.-hydroxylation** product was identified by gas chromatog. on a polyester column and reverse phase, high performance liq. chromatog., 2 techniques which were shown to resolve the positional isomers. Gas chromatog. anal. of the **dicarboxylic acid** obtained by CrO₃ oxidn. of the enzymic product also confirmed the identity

of the enzymic **.omega.-hydroxylation** product. This enzymic **hydroxylation** required O and NADPH, but substitution of NADH resulted in nearly half the reaction rate obtained with NADPH. Maximal rates of **.omega.-hydroxylation** occurred at pH 8 and the rate increased in a sigmoidal manner with increasing concns. of palmitic acid. This **.omega.-hydroxylation** was inhibited by the classical **mixed function oxidase** inhibitors such as metal chelators (o-phenanthroline, 8-hydroxyquinoline, and **.alpha.,.alpha.-dipyridyl**), NaN₃, and thiol reagents (N-ethylmaleimide and p-chloromercuribenzoate). As expected of a hydroxylase, involving cytochrome P450, the present o-hydroxylase was inhibited by CO and this enzyme system showed unusually high sensitivity to this inhibition; 10% CO caused inhibition and 30% CO completely inhibited the reaction. Another unusual feature was that the inhibition caused by any level of CO could not be reversed by light (420-460 nm).

TI Biosynthesis of cutin. **.omega.-Hydroxylation** of fatty acids by a microsomal **preparation** from germinating *Vicia faba*

AB **.omega.-Hydroxylation** of fatty acids, which is a key reaction in the biosynthesis of cutin and suberin, was demonstrated in a cell-free **prepn.** from a higher plant. A crude microsomal fraction (105,000-g pellet) from germinating embryonic shoots of *V. faba* catalyzed the conversion of palmitic acid to **.omega.-hydroxypalmitic**

acid.

As the crude cell-free **prepn.** also catalyzes the formation of other hydroxy acids such as **.alpha.-** and **.beta.-hydroxy** acids, the **.omega.-hydroxylation** product was identified by gas chromatog. on a polyester column and reverse phase, high performance liq. chromatog., 2 techniques which were shown to resolve the positional isomers. Gas chromatog. anal. of the **dicarboxylic acid** obtained by CrO₃ oxidn. of the enzymic product also confirmed the identity

of the enzymic **.omega.-hydroxylation** product. This enzymic **hydroxylation** required O and NADPH, but substitution of NADH resulted in nearly half the reaction rate obtained with NADPH. Maximal rates of **.omega.-hydroxylation** occurred at pH 8 and the rate increased in a sigmoidal manner with increasing concns. of palmitic acid. This **.omega.-hydroxylation** was inhibited by the classical **mixed function oxidase** inhibitors such as metal chelators (o-phenanthroline, 8-hydroxyquinoline, and **.alpha.,.alpha.-dipyridyl**), NaN₃, and thiol reagents (N-ethylmaleimide and p-chloromercuribenzoate). As expected of a hydroxylase, involving cytochrome P450, the present o-hydroxylase was inhibited by CO and this enzyme system showed unusually

high sensitivity to this inhibition; 10% CO caused inhibition and 30% CO completely inhibited the reaction. Another unusual feature was that the inhibition caused by any level of CO could not be reversed by light (420-460 nm).

ST **omega hydroxylation** fatty acid Vicia

IT Fatty acids, biological studies

RL: RCT (Reactant)

(**.omega.-hydroxylation** of, by microsome from broad bean)

IT Hydroxylation

(omega, of fatty acids by microsomal **prepn.** from Vicia faba)

=> d full history

(FILE 'HOME' ENTERED AT 08:53:03 ON 10 MAY 2001)

FILE 'CAPLUS' ENTERED AT 08:53:11 ON 10 MAY 2001

E .BETA.-OXIDN.-BLOCKED/CT
E E2
E E3+ALL
E .BETA.-OXIDATION/CT
E E3+ALL

FILE 'REGISTRY' ENTERED AT 09:37:48 ON 10 MAY 2001

L1 1 SEA ABB=ON PLU=ON 9038-14-6/RN
D
L2 1 SEA ABB=ON PLU=ON 9023-03-4/RN
D

FILE 'CAOLD, CAPLUS, CROPU, DGENE, DPCI, ENCOMPPAT, ENCOMPPAT2, EUROPATFULL, IFIPAT, INPADOC, JAPIO, PAPERCHEM2, PATDD, PATDPA, PATOSDE, PATOSEP, PATOSWO, PCTFULL, PIRA, RAPRA, SYNTHLINE, TULSA, TULSA2, USPATFULL, WPIDS' ENTERED AT 09:39:31 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 09:39:57 ON 10 MAY 2001

L3 SET SMARTSELECT ON
SEL PLU=ON L1 1- CHEM : 26 TERMS
SET SMARTSELECT OFF

FILE 'CAOLD, CAPLUS, CROPU, DGENE, DPCI, ENCOMPPAT, ENCOMPPAT2, EUROPATFULL, IFIPAT, INPADOC, JAPIO, PAPERCHEM2, PATDD, PATDPA, PATOSDE, PATOSEP, PATOSWO, PCTFULL, PIRA, RAPRA, SYNTHLINE, TULSA, TULSA2, USPATFULL, WPIDS' ENTERED AT 09:40:22 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 09:40:25 ON 10 MAY 2001

L4 SET SMARTSELECT ON
SEL PLU=ON L2 1- CHEM : 16 TERMS
SET SMARTSELECT OFF

FILE 'CAOLD, CAPLUS, CROPU, DGENE, DPCI, ENCOMPPAT, ENCOMPPAT2, EUROPATFULL, IFIPAT, INPADOC, JAPIO, PAPERCHEM2, PATDD, PATDPA, PATOSDE, PATOSEP, PATOSWO, PCTFULL, PIRA, RAPRA, SYNTHLINE, TULSA, TULSA2, USPATFULL, WPIDS' ENTERED AT 09:40:33 ON 10 MAY 2001

L5 18788 SEA ABB=ON PLU=ON L3
L6 7868 SEA ABB=ON PLU=ON L4
L7 25412 SEA ABB=ON PLU=ON L5 OR L6
L8 830900 SEA ABB=ON PLU=ON DICARBOXYLIC ACID# OR (CARBOXYLIC ACIDS
(L) DICARBOXYLIC) OR CARBOXYLIC ACID# OR MONOCARBOXYLIC ACID#
L9 573 SEA ABB=ON PLU=ON L7 AND L8
L10 361 SEA ABB=ON PLU=ON L9 AND (PREPAR? OR SYNTHES? OR MAK? OR
PREP/RL)
L11 11119 SEA ABB=ON PLU=ON (HYDROXYLATION (L) .OMEGA.-HYDROXYLATION)
OR (HYDROXYLATION (L) ALKANE#) OR (.ALPHA.-HYDROXYLATION) OR
(HYDROXYLATION (L) .ALPHA.)
L12 76 SEA ABB=ON PLU=ON L11 AND L10
L13 74 DUP REM L12 (2 DUPLICATES REMOVED)
L14 1066 SEA ABB=ON PLU=ON CANDIDA MALTOSA OR CANDIDA CLOACAE OR
CANDIDA NOVELLUS OR CANDIDA SUBTROPICALIS
L15 7 SEA ABB=ON PLU=ON L14 AND L10
L16 6 DUP REM L15 (1 DUPLICATE REMOVED)
D IBIB AB 1-6

D IBIB AB 1
D IBIB AB 1-6

=> d ibib ab 1-6

L16 ANSWER 1 OF 6 PCTFULL COPYRIGHT 2001 MicroPatent
ACCESSION NUMBER: 2000065061 PCTFULL EW 200044 ED 20001124
TITLE (ENGLISH): NUCLEIC ACID SEQUENCES FROM <i> CANDIDA </i> YEASTS
WHICH CODE
CYTOCHROME B5 POLYPEPTIDES
TITLE (FRENCH): SEQUENCES D'ACIDE NUCLEIQUE ISSUES DE LEVURES DE <i>
CANDIDA </i>,
CODANT DES POLYPEPTIDES B5 CYTOCHROMES
TITLE (GERMAN): NUKLEINSAEURE-SEQUENZEN AUS <i> CANDIDA </i> HEFEN,
DIE CYTOCHROM
B5-POLYPEPTIDE KODIEREN
INVENTOR(S): SCHUNCK, Wolf-Hagen; CHERNOGOLOV, Alexei
PATENT ASSIGNEE(S): MAX-DELB RUECK-CENTRUM FUE R MOLEKULARE MEDIZIN
LANGUAGE OF PUBL.: German
LANGUAGE OF FILING: German
DOCUMENT TYPE: Patent
PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 2000065061	A2	20001102
DESIGNATED STATES:	AE AG AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DK DM DZ EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MA MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT TZ UA UG US UZ VN YU ZA ZW GH GM KE LS MW SD SL SZ TZ UG ZW AM AZ BY KG KZ MD RU TJ TM AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN GW ML MR NE SN TD TG		
APPLICATION INFO.:	WO 2000-DE1246		20000418
PRIORITY (ORIGINAL):	DE 1999-199 18 763.0		19990424
ABEN	The invention relates to nucleic acid sequences from <i> Candida </i> yeasts, preferably from <i> Candida maltosa </i>, which code cytochrome b5 polypeptides, as well as to the corresponding cytochrome b5 polypeptides and their use for increasing the activity of cytochrome P450 systems, especially for stimulating the activity of alkane- hydroxylating and fatty acid-hydroxylating cytochrome P450 systems during the production of long-chained dicarboxylic acids (#ge#C10).		
ABFR	L'invention concerne des sequences d'acide nucleique issues de levures de <i> Candida </i>, de preference de <i> Candida maltosa </i>, qui codent des polypeptides b5 cytochromes. L'invention concerne egalement les polypeptides b5 cytochromes correspondants et leur utilisation pour augmenter l'activite de systemes P450 cytochromes, notamment pour stimuler l'activite de systemes P450 cytochromes a effet hydroxylant de l'alcane et de l'acide gras lors de la preparation d'acides dicarboxyliques a chaine longue (#ge#C10).		
ABDE	Die Erfindung betrifft Nukleinsaeure-Sequenzen aus <i> Candida </i> Hefen vorzugsweise aus <i> Candida maltosa </i>, die Cytochrom b5- Polypeptide kodieren sowie die entsprechenden Cytochrom b5-Polypeptide und ihre Verwendung zur Erhoehung der Aktivitaet von Cytochrom P450 Systemen, insbesondere zur Stimulierung der Aktivitaet Alkan- und Fettsaeure-hydroxylierender Cytochrom P450 Systeme bei der Produktion langkettiger Dicarbonsaeuren (#ge#C10).		

L16 ANSWER 2 OF 6
ACCESSION NUMBER:
TITLE (ENGLISH):
TITLE (FRENCH):
INVENTOR(S):
PATENT ASSIGNEE(S):
LANGUAGE OF PUBL.:
LANGUAGE OF FILING:
DOCUMENT TYPE:
PATENT INFORMATION:

PCTFULL COPYRIGHT 2001 MicroPatent
2000034473 PCTFULL EW 200024 ED 20000712
SEVEN TRANSMEMBRANE DOMAIN RECEPTOR ZSIG56
DOMAINE TRANSMEMBRANAIRE 7 ZSIG56
SHEPPARD, Paul, O.; ELLSWORTH, Jeff, L.
ZYMOGENETICS, INC.
English
English
Patent

	NUMBER	KIND	DATE
	WO 2000034473	A2	20000615
DESIGNATED STATES:	AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CR CU CZ DE DK DM EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA UG UZ VN YU ZA ZW GH GM KE LS MW SD SL SZ TZ UG ZW AM AZ BY KG KZ MD RU TJ TM AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN GW ML MR NE SN TD TG		
APPLICATION INFO.:	WO 1999-US28492		19991202
PRIORITY (ORIGINAL):	US 1998-09/208691		19981210
ABEN	The present invention relates to polynucleotide and polypeptide molecules for a seven transmembrane domain receptor designated zsig56. The polypeptides, and polynucleotides encoding them are useful treating pathological conditions in such diverse tissue, as kidney, thyroid, gastrointestinal, CSN and reproductive. Such conditions include hypertension, hyper- and hypothyroidism, neurotransmission, gastrointestinal motility, inflammation and reproduction. The present invention also includes antibodies to the zsig56 polypeptides.		
ABFR	L'invention concerne des molecules polynucleotidiques et polypeptidiques pour un recepteur du domaine transmembranaire 7 appele zsig56. Les polypeptides et les polynucleotides codant pour eux conviennent pour le traitement d'etats pathologiques de tissus aussi divers que les reins, la glande thyroide, le tube digestif, le systeme nerveux central et l'appareil genital. Lesdits etats pathologiques concernent l'hypertension, l'hyperthyroidie, l'hypothyroidie, la neurotransmission, le transit intestinal, l'inflammation et la reproduction. L'invention concerne egalement des anticorps diriges contre les polypeptides zsig56.		

L16 ANSWER 3 OF 6
ACCESSION NUMBER:
TITLE (ENGLISH):

PCTFULL COPYRIGHT 2001 MicroPatent
2000020566 PCTFULL EW 200015 ED 20000503
CYTOCHROME P450 **MONOOXYGENASE** AND NADPH
CYTOCHROME P450
OXIDOREDUCTASE GENES AND PROTEINS RELATED TO THE

OMEGA

HYDROXYLASE
COMPLEX OF *CANDIDA TROPICALIS* AND METHODS
RELATING THERETO

TITLE (FRENCH):

GENES DE LA **MONOOXYGENASE** DU CYTOCHROME P450
ET DE
L'OXYDOREDUCTASE NADPH DU CYTOCHROME P450 ET

PROTEINES

ASSOCIEES AU
COMPLEXE DE L'OMEGA HYDROXYLASE DE *CANDIDA TROPICALIS* ET PROCEDES
ASSOCIES

INVENTOR(S):
Dudley;

WILSON, C., Ron; CRAFT, David, L.; EIRICH, L.,

A.; ESHOO, Mark; MADDURI, Krishna, M.; CORNETT, Cathy,
 BRENNER, Alfred, A.; TANG, Maria; LOPER, John, C.;
 GLEESON, Martin
 PATENT ASSIGNEE(S): HENKEL CORPORATION
 LANGUAGE OF PUBL.: English
 LANGUAGE OF FILING: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 2000020566	A2	20000413
DESIGNATED STATES:	AE AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GD GE GH GM HR HU ID IL IN IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK SL TJ TM TR TT UA UG US UZ VN YU ZA ZW GH GM KE LS MW SD SL SZ UG ZW AM AZ BY KG KZ MD RU TJ TM AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN GW ML MR NE SN TD TG		
APPLICATION INFO.:	WO 1999-US20797		19990910
PRIORITY (ORIGINAL):	US 1998-60/103099		19981005
	US 1999-		19990310

ABEN Novel genes have been isolated which encode cytochrome P450 and NADPH reductase enzyme of the #ohgr#-hydroxylase complex of *C. tropicalis* 20336. Vectors including these genes, transfected host cells and transformed host cells are provided. Methods of producing of cytochrome P450 and NADPH reductase enzymes are also provided which involve transforming a host cell with a gene encoding these enzymes and culturing the cells. Methods of increasing the production of a **dicarboxylic acid** and methods of increasing production of the aforementioned enzymes are also provided which involve increasing in the host cell the number of genes encoding these enzymes. A method for discriminating members of a gene family by quantifying the expression of genes is also provided.

ABFR La presente invention concerne des genes isolees qui codent pour des enzymes du cytochrome P450 et de reductase de NADPH du complexe #ohgr#-hydroxylase de *C. tropicalis* 20336. L'invention concerne aussi des vecteurs contenant ces genes, des cellules hotes transfectees ainsi que des cellules hotes transformees. Sont aussi decrits des procedes de production des enzymes du cytochrome P450 et de reductase de NADPH qui consistent a transformer une cellule hote a l'aide d'un gene qui code pour ces enzymes et a cultiver les cellules. L'invention concerne aussi des procedes d'augmentation de la production d'un acide dicarboxylique, des procedes d'augmentation de la production des enzymes mentionnees ci-dessus qui consistent a augmenter le nombre de genes qui codent pour ces enzymes dans les cellules hotes, ainsi qu'une methode destinee a distinguer des elements d'une famille de genes en quantifiant l'expression des genes.

L16 ANSWER 4 OF 6 CAPLUS COPYRIGHT 2001 ACS DUPLICATE 1
 ACCESSION NUMBER: 1999:77701 CAPLUS
 DOCUMENT NUMBER: 130:138397
 TITLE: Yeasts with elevated cytochrome P450 levels and their use in the manufacture of monoterminial and diterminial aliphatic carboxylates from alkanes
 INVENTOR(S): Fallon, Robert D.; Payne, Mark S.; Picataggio, Stephen
 K.; Wu, Shijun

PATENT ASSIGNEE(S): E.I. Du Pont De Nemours and Company, USA
 SOURCE: PCT Int. Appl., 58 pp.
 CODEN: PIXXD2
 DOCUMENT TYPE: Patent
 LANGUAGE: English
 FAMILY ACC. NUM. COUNT: 1
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9904014	A2	19990128	WO 1998-US14935	19980720
WO 9904014	A3	19990520		
W: AU, CA, IL, JP, NZ, US				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9884982	A1	19990210	AU 1998-84982	19980720
EP 1003881	A2	20000531	EP 1998-935806	19980720
R: DE, DK, ES, FR, GB, IT, NL, SE, IE				

PRIORITY APPLN. INFO.: US 1997-53215 P 19970721
 WO 1998-US14935 W 19980720

AB Strains of *Pichia pastoris* and *Candida maltosa* that have increased cytochrome P 450 activity or lack the .beta.-oxidn. pathway that can be used to convert c6-C22 alkanes to monoterminial and diterminial **carboxylic acids** are described. Expression constructs carrying cytochrome P 450 genes under the control of powerful inducible promoters (AOX1, PGK) were constructed by std. methods and introduced into *P. pastoris* or *C. maltosa* to give novel or increased P 450 activities. Disruption of .beta.-oxidn. in *C. maltosa* was achieved by insertional inactivation of the POX4 gene. A strain of *C. maltosa* cultured in a complete medium that was supplemented with dodecane 20 g/L when glucose was almost completely depleted was able to generate dodecanedioic acid at 3.4 g/h for 51 h.

L16 ANSWER 5 OF 6 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1998055612 PCTFULL
 TITLE (ENGLISH): NEUROKININ B PRECURSORS
 TITLE (FRENCH): PRECURSEURS DE NEUROKININE B
 INVENTOR(S): SHEPPARD, Paul, O.
 PATENT ASSIGNEE(S): ZYMOGENETICS, INC.
 LANGUAGE OF PUBL.: English
 LANGUAGE OF FILING: English
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
	WO 9855612	A1	19981210
DESIGNATED STATES:	AL AM AT AU AZ BA BB BG BR BY CA CH CN CU CZ DE DK EE ES FI GB GE GH HU IL IS JP KE KG KP KR KZ LC LK LR LS LT LU LV MD MG MK MN MW MX NO NZ PL PT RO RU SD SE SG SI SK TJ TM TR TT UA UG UZ VN ZW GH GM KE LS MW SD SZ UG ZW AM AZ BY KG KZ MD RU TJ TM AT BE CH CY DE DK ES FI FR GB GR IE IT LU MC NL PT SE BF BJ CF CG CI CM GA GN ML MR NE SN TD TG		
APPLICATION INFO.:	WO 1998-US10842		19980528
PRIORITY (ORIGINAL):	US 1997-60/048290		19970602

ABEN The present invention relates to zneurokl polypeptides and polynucleotides encoding the same. These polypeptides are novel members of a family of proteins that are precursors of neurokinin B, a ten amino

acid moiety of biological significance. The polypeptides, and polynucleotides encoding them, are useful in the study of prohormone convertase function and neurokinin receptors. The present invention also includes antibodies to the zneurokl polypeptides.

ABFR Cette invention concerne des polypeptides zneurokl ainsi que des polynucleotides codant ces derniers. Ces polypeptides consistent en de nouveaux membres d'une famille de proteines qui sont des precurseurs de neurokinine B, un fragment de l'acide amine dix jouant un role biologique important. Ces polypeptides, ainsi que les polynucleotides qui les codent, sont utiles lors de l'etude de la fonction de convertase de prohormone, ainsi que des recepteurs de neurokinine. Cette invention concerne egalement des anticorps diriges contre les polypeptides zneurokl.

L16 ANSWER 6 OF 6 PCTFULL COPYRIGHT 2001 MicroPatent
 ACCESSION NUMBER: 1996027678 PCTFULL
 TITLE (ENGLISH): PROCESS FOR HYDROXYLATING LONG-CHAIN ALKANES, FATTY ACIDS AND OTHER ALKYL COMPOUNDS
 TITLE (FRENCH): PROCEDE D'HYDROXYLATION D'ALCANES A LONGUE CHAINE, D'ACIDES GRAS ET D'AUTRES COMPOSES ALKYLE
 INVENTOR(S): ZIMMER, Thomas; KAMINSKI, Kristina; SCHUNCK, Wolf-Hagen; KAERGEL, Eva; SCHELLER, Ulrich; MAUERSBERGER, Stephan
 PATENT ASSIGNEE(S): MAX-DELBROECK-CENTRUM FUER MOLEKULARE MEDIZIN; ZIMMER, Thomas; KAMINSKI, Kristina; SCHUNCK, Wolf-Hagen; KAERGEL, Eva; SCHELLER, Ulrich; MAUERSBERGER, Stephan
 LANGUAGE OF PUBL.: German
 DOCUMENT TYPE: Patent
 PATENT INFORMATION:

	NUMBER	KIND	DATE
DESIGNATED STATES:	WO 9627678	A1	19960912
	JP US AT BE CH DE DK ES FI FR GB GR IE IT LU MC NL PT SE		
APPLICATION INFO.:	WO 1996-DE410		19960301
PRIORITY (ORIGINAL):	DE 1995-195 07 546.3		19950303

ABEN A microbial hydroxylation process is disclosed for selectively oxidising regions of long-chain alkanes, fatty acids and other alkyl compounds. The process should be easy to carry out and produce good yields of oxidation products, in particular hydroxylated fatty acids and long-chain **dicarboxylic acids**. The object of the invention is to modify yeast by genetic engineering so that when it is cultivated it expresses the required enzymes. The disclosed process is characterised in that the long-chain alkanes, fatty acids and other alkyl compounds are treated with **monooxygenase** systems that consist of cytochrom P450 and NADPOH cytochrom P450 reductase, and the hydroxylation products are then separated. The **monooxygenase** systems are produced in the reaction mixture by simultaneous expression of their components in yeast, preferably *saccharomyces cerevisiae*. The invention relates essentially to a vector for modifying *saccharomyces* by genetic engineering. On the basis of the structure Yep 51, the vector contains reductase cDNA between the restriction sites SalK and BamHI and a second expression cassette bound in the restriction site NruI. The expression cassette consists of the GAL10 promoter, the sequence that codes for cytochrom

P450 and the ADH1 terminator.

ABF L'invention concerne un procede d'hydroxylation microbienne qui sert a oxyder selectivement des regions d'alcanes a longue chaine, d'acides gras et d'autres composes alkyle. Le procede doit permettre d'obtenir de maniere aisee un bon rendement en produits d'oxydation, notamment des acides gras'hydroxyles et des acides dicarboxyliques a longue chaine. L'invention a pour objet de modifier par genie genetique des levures de sorte qu'elles expriment les enzymes requises lorsqu'elles sont cultivees. Le procede se caracterise en ce que l'on traite les alcanes a longue chaine, les acides gras et les autres composes alkyle avec des systemes a **monooxygenase** constitues de cytochrome P450 et de NADPH-cytochrome P450-reductase, les produits d'hydroxylation etant ensuite isoles. Les systemes a **monooxygenase** sont produits dans le melange de reaction par expression simultanee de leurs composants dans des levures, de preference la *saccharomyces cerevisiae*. L'invention porte essentiellement sur un vecteur de modification par genie genetique de *saccharomyces*, qui sur la base de la structure fondamentale Yep 51 contient l'ADNc de la reductase entre les sites de restriction SalI et BamHI et une deuxieme cassette d'expression liee dans le site de restriction NruI et constituee du promoteur GAL10, de la sequence de codage de cytochrome P450 et du terminateur ADH1.

=> d full history

(FILE 'HOME' ENTERED AT 11:17:32 ON 10 MAY 2001)

FILE 'REGISTRY' ENTERED AT 11:17:52 ON 10 MAY 2001
L1 1 SEA ABB=ON PLU=ON 9038-14-6/RN
D
L2 1 SEA ABB=ON PLU=ON 9023-03-4/RN
D

FILE 'HCAPLUS' ENTERED AT 11:18:36 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 11:18:51 ON 10 MAY 2001
SET SMARTSELECT ON
L3 SEL PLU=ON L1 1- CHEM : 26 TERMS
SET SMARTSELECT OFF

FILE 'HCAPLUS' ENTERED AT 11:18:52 ON 10 MAY 2001

FILE 'REGISTRY' ENTERED AT 11:18:53 ON 10 MAY 2001
SET SMARTSELECT ON
L4 SEL PLU=ON L2 1- CHEM : 16 TERMS
SET SMARTSELECT OFF

FILE 'HCAPLUS' ENTERED AT 11:18:54 ON 10 MAY 2001
L5 15271 SEA ABB=ON PLU=ON L3
L6 7478 SEA ABB=ON PLU=ON L4
L7 21584 SEA ABB=ON PLU=ON L5 OR L6
L8 188643 SEA ABB=ON PLU=ON DICARBOXYLIC ACID# OR (CARBOXYLIC ACIDS
(L) DICARBOXYLIC) OR CARBOXYLIC ACID# OR MONOCARBOXYLIC ACID#
L9 5883 SEA ABB=ON PLU=ON (HYDROXYLATION (L) .OMEGA.-HYDROXYLATION)
OR (HYDROXYLATION (L) ALKANE#) OR (.ALPHA.-HYDROXYLATION) OR
(HYDROXYLATION (L) .ALPHA.)
L10 1338 SEA ABB=ON PLU=ON PICHIA PASTORIS
L11 4 SEA ABB=ON PLU=ON L7 (L) L8 (L) L9
D IBIB AB 1-4
D IBIB AB HIT 1-4
L12 0 SEA ABB=ON PLU=ON L11 AND L10

=> d ibib ab 1-4

L11 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1993:5685 HCAPLUS

DOCUMENT NUMBER: 118:5685

TITLE: Metabolic engineering of *Candida tropicalis* for the production of long-chain dicarboxylic acids

AUTHOR(S): Picataggio, Stephen; Rohrer, Tracy; Deanda, Kristine; Lanning, Dawn; Reynolds, Robert; Mielenz, Jonathan; Eirich, L. Dudley

CORPORATE SOURCE: Microb. Technol. Dep., Cognis Inc., Santa Rosa, CA, 95407, USA

SOURCE: Bio/Technology (1992), 10(8), 894-8

CODEN: BTCHDA; ISSN: 0733-222X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An industrial strain of the yeast *C. tropicalis* was engineered for the efficient prodn. of long-chain **dicarboxylic acids**, which are important raw materials for the chem. industry. By sequential disruption of the 4 genes encoding both isoenzymes of the acyl-CoA oxidase

which catalyzes the first reaction in the .beta.-oxidn. pathway, **alkane** and fatty acid substrates were successfully redirected to the .omega.-oxidn. pathway. Consequently, the conversion efficiency and chem. selectivity of their terminal oxidn. to the corresponding **dicarboxylic acids** was improved to 100%. The specific productivity of the bioconversion was increased further by amplification of the **cytochrome P 450 monooxygenase** and NADPH-cytochrome reductase genes encoding the rate-limiting .omega.-hydroxylase in the .omega.-oxidn. pathway. The amplified strains demonstrated increased .omega.-hydroxylase activity and a 30% increase in productivity compared to the .beta.-oxidn.-blocked strain in fermns. The bioconversion is effective for the selective terminal oxidn. of both satd. and unsatd. linear aliph. substrates with C12-22 chain-lengths and also avoids the undesirable chain modifications assocd. with passage through the .beta.-oxidn. pathway, such as unsatn., **hydroxylation**, or chain shortening. It is now possible to efficiently produce a wide range of previously unavailable satd. and unsatd. **dicarboxylic acids** with a high degree of purity.

L11 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1991:220705 HCAPLUS

DOCUMENT NUMBER: 114:220705

TITLE: Microsomal oxidation of dodecylthioacetic acid (a 3-thia fatty acid) in rat liver

AUTHOR(S): Hvattum, Erlend; Bergseth, Steinar; Pedersen, Catharina N.; Bremer, Jon; Aarsland, Asle; Berge, Rolf

K.

CORPORATE SOURCE: Inst. Med. Biochem., Univ. Oslo, Oslo, 0317, Norway

SOURCE: Biochem. Pharmacol. (1991), 41(6-7), 945-53

CODEN: BCPCA6; ISSN: 0006-2952

DOCUMENT TYPE: Journal

LANGUAGE: English

AB [1-14C]Dodecylthioacetic acid (DTA), a 3-thia fatty acid, is .omega. (.omega.-1)-hydroxylated and sulfur oxygenated at about equal rates in rat

liver microsomes. In prolonged incubations DTA is converted to

.omega.-hydroxydodecylsulfoxyacetic acid. .omega.-
Hydroxylation of DTA is catalyzed by cytochrome P450IVA1 (or a
 very closely related isoenzyme in the same gene family), the fatty acid
 .omega.-hydroxylating enzyme. It is absolutely dependent on NADPH and
 inhibited by CO, and lauric acid is a competing substrate. .omega.
 .-**Hydroxylation** of DTA is increased by feeding
 tetradecylthioacetic acid (TTA), a 3-thia fatty acid, for 4 days to rats.
 .omega.-**Hydroxylation** of [1-14C] lauric acid is also
 induced by TTA and other 3-thia **carboxylic acids**. A
 close relationship was obsd. between induction of microsomal .
omega.-hydroxylation of fatty acid and palmitoyl-CoA
 hydrolase activity. DTA is .omega.-hydroxylated at about the same rate

as

the physiol. substrate lauric acid. The sulfur oxygenation of DTA is
 catalyzed by liver microsomal **flavin-contg.**
monooxygenase (FMO) (EC 1.14.13.8). It is dependent on either
 NADH or NADPH. The Km value for NADH was .apprx.five times larger than
 the Km value for NADPH. It is inhibited by methimazole and not affected
 by CO. It is not induced by TTA.

L11 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1985:61628 HCAPLUS
 DOCUMENT NUMBER: 102:61628
 TITLE: Oxidative ether cleavage with p-nitroperbenzoic acid
 AUTHOR(S): Schneider, Hans Joerg; Ahlhelm, Alfred; Mueller,
 Walter
 CORPORATE SOURCE: Univ. Saarlandes, Saarbruecken, D-6600/11, Fed. Rep.
 Ger.
 SOURCE: Chem. Ber. (1984), 117(11), 3297-302
 CODEN: CHBEAM; ISSN: 0009-2940
 DOCUMENT TYPE: Journal
 LANGUAGE: German

AB The reaction of p-nitroperbenzoic acid in CHCl3 with Me2CH(CH2)3OMe and
 (Me2CHCH2CH2)2O leads by selective attack at C-H bonds in the .
alpha.-position to the ether oxygen to hemiacetals, which decomp.
 to aldehydes and alcs., yielding **carboxylic acids**.
 Secondary alkoxy groups, as in ethoxycyclohexane, furnish Baeyer-Villiger
 oxidn. products of initially formed ketones. Kinetic measurements with
 substituted benzyl Me ethers show a Hammett reaction const. .rho. = -0.9,
 which is in accord with the obsd. relatively small discrimination between
 secondary and tertiary C-H bonds. The results are compared with similar
hydroxylations of **alkanes** and with **monooxygenase**
 reactions and point to oxenoid transition states. Radical reactions, as
 found with some **alkanes** are not obsd., which is shown by the
 small amts. of PhNO2 (.ltoreq.10%) formed during the reaction. 13C-NMR
 shifts of several ethers and oxidn. products are reported.

L11 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1977:498893 HCAPLUS
 DOCUMENT NUMBER: 87:98893
 TITLE: Biosynthesis of cutin. .omega.-Hydroxylation of
 fatty
 acids by a microsomal preparation from germinating
 Vicia faba
 AUTHOR(S): Soliday, Charles L.; Kolattukudy, P. E.
 CORPORATE SOURCE: Dep. Agric. Chem., Washington State Univ., Pullman,
 Wash., USA
 SOURCE: Plant Physiol. (1977), 59(6), 1116-21
 CODEN: PLPHAY
 DOCUMENT TYPE: Journal

LANGUAGE: English

AB **.omega.-Hydroxylation** of fatty acids, which is a key reaction in the biosynthesis of cutin and suberin, was demonstrated in a cell-free prepn. from a higher plant. A crude microsomal fraction (105,000-g pellet) from germinating embryonic shoots of *V. faba* catalyzed the conversion of palmitic acid to **.omega.-hydroxypalmitic acid**. As the crude cell-free prepn. also catalyzes the formation of other hydroxy

acids

such as **.alpha.-** and **.beta.-hydroxy acids**, the **.omega.-hydroxylation** product was identified by gas chromatog. on a polyester column and reverse phase, high performance liq. chromatog., 2 techniques which were shown to resolve the positional isomers. Gas chromatog. anal. of the **dicarboxylic acid** obtained by CrO₃ oxidn. of the enzymic product also confirmed the identity of the enzymic **.omega.-hydroxylation** product. This enzymic **hydroxylation** required O and NADPH, but substitution of NADH resulted in nearly half the reaction rate obtained with NADPH. Maximal rates of **.omega.-hydroxylation** occurred at pH 8 and the rate increased in a sigmoidal manner with increasing concns. of palmitic acid. This **.omega.-hydroxylation** was inhibited by the classical **mixed function oxidase** inhibitors such as metal chelators (o-phenanthroline, 8-hydroxyquinoline, and **.alpha.,.alpha.-dipyridyl**), NaN₃, and thiol reagents (N-ethylmaleimide and p-chloromercuribenzoate). As expected of a hydroxylase, involving cytochrome P450, the present o-hydroxylase was inhibited by CO and this enzyme system showed unusually high sensitivity to this inhibition; 10% CO caused inhibition and 30% CO completely inhibited the reaction. Another unusual feature was that the inhibition caused by any level of CO could not be reversed by light (420-460 nm).

=> d ibib ab hit 1-4

L11 ANSWER 1 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1993:5685 HCAPLUS

DOCUMENT NUMBER: 118:5685

TITLE: Metabolic engineering of *Candida tropicalis* for the production of long-chain dicarboxylic acids

AUTHOR(S): Picataggio, Stephen; Rohrer, Tracy; Deanda, Kristine; Lanning, Dawn; Reynolds, Robert; Mielenz, Jonathan; Eirich, L. Dudley

CORPORATE SOURCE: Microb. Technol. Dep., Cognis Inc., Santa Rosa, CA, 95407, USA

SOURCE: Bio/Technology (1992), 10(8), 894-8

CODEN: BTCHDA; ISSN: 0733-222X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An industrial strain of the yeast *C. tropicalis* was engineered for the efficient prodn. of long-chain **dicarboxylic acids**, which are important raw materials for the chem. industry. By sequential disruption of the 4 genes encoding both isoenzymes of the acyl-CoA oxidase

which catalyzes the first reaction in the **.beta.-oxidn. pathway**, **alkane** and fatty acid substrates were successfully redirected to the **.omega.-oxidn. pathway**. Consequently, the conversion efficiency and chem. selectivity of their terminal oxidn. to the corresponding **dicarboxylic acids** was improved to 100%. The specific productivity of the bioconversion was increased further by amplification of the **cytochrome P 450**

monooxygenase and NADPH-cytochrome reductase genes encoding the rate-limiting .omega.-hydroxylase in the .omega.-oxidn. pathway. The amplified strains demonstrated increased .omega.-hydroxylase activity and a 30% increase in productivity compared to the .beta.-oxidn.-blocked strain in fermns. The bioconversion is effective for the selective terminal oxidn. of both satd. and unsatd. linear aliph. substrates with C12-22 chain-lengths and also avoids the undesirable chain modifications assocd. with passage through the .beta.-oxidn. pathway, such as unsatn., **hydroxylation**, or chain shortening. It is now possible to efficiently produce a wide range of previously unavailable satd. and unsatd. **dicarboxylic acids** with a high degree of purity.

AB An industrial strain of the yeast *C. tropicalis* was engineered for the efficient prodn. of long-chain **dicarboxylic acids**, which are important raw materials for the chem. industry. By sequential disruption of the 4 genes encoding both isoenzymes of the acyl-CoA oxidase

which catalyzes the first reaction in the .beta.-oxidn. pathway, **alkane** and fatty acid substrates were successfully redirected to the .omega.-oxidn. pathway. Consequently, the conversion efficiency and chem. selectivity of their terminal oxidn. to the corresponding **dicarboxylic acids** was improved to 100%. The specific productivity of the bioconversion was increased further by amplification of the **cytochrome P 450**

monooxygenase and NADPH-cytochrome reductase genes encoding the rate-limiting .omega.-hydroxylase in the .omega.-oxidn. pathway. The amplified strains demonstrated increased .omega.-hydroxylase activity and a 30% increase in productivity compared to the .beta.-oxidn.-blocked strain in fermns. The bioconversion is effective for the selective terminal oxidn. of both satd. and unsatd. linear aliph. substrates with C12-22 chain-lengths and also avoids the undesirable chain modifications assocd. with passage through the .beta.-oxidn. pathway, such as unsatn., **hydroxylation**, or chain shortening. It is now possible to efficiently produce a wide range of previously unavailable satd. and unsatd. **dicarboxylic acids** with a high degree of purity.

L11 ANSWER 2 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1991:220705 HCAPLUS

DOCUMENT NUMBER: 114:220705

TITLE: Microsomal oxidation of dodecylthioacetic acid (a 3-thia fatty acid) in rat liver

AUTHOR(S): Hvattum, Erlend; Bergseth, Steinar; Pedersen, Catharina N.; Bremer, Jon; Aarsland, Asle; Berge,

Rolf

K.

CORPORATE SOURCE: Inst. Med. Biochem., Univ. Oslo, Oslo, 0317, Norway

SOURCE: Biochem. Pharmacol. (1991), 41(6-7), 945-53

CODEN: BCPCA6; ISSN: 0006-2952

DOCUMENT TYPE: Journal

LANGUAGE: English

AB [1-14C]Dodecylthioacetic acid (DTA), a 3-thia fatty acid, is .omega. (.omega.-1)-hydroxylated and sulfur oxygenated at about equal rates in rat

liver microsomes. In prolonged incubations DTA is converted to .omega.-hydroxydodecylsulfoxyacetic acid. .omega.-

Hydroxylation of DTA is catalyzed by cytochrome P450IVA1 (or a very closely related isoenzyme in the same gene family), the fatty acid .omega.-hydroxylating enzyme. It is absolutely dependent on NADPH and inhibited by CO, and lauric acid is a competing substrate. .omega.

.-Hydroxylation of DTA is increased by feeding tetradecylthioacetic acid (TTA), a 3-thia fatty acid, for 4 days to rats. .omega.-Hydroxylation of [1-14C] lauric acid is also induced by TTA and other 3-thia **carboxylic acids**. A close relationship was obsd. between induction of microsomal .omega.-hydroxylation of fatty acid and palmitoyl-CoA hydrolase activity. DTA is .omega.-hydroxylated at about the same rate as the physiol. substrate lauric acid. The sulfur oxygenation of DTA is catalyzed by liver microsomal **flavin-contg. monooxygenase** (FMO) (EC 1.14.13.8). It is dependent on either NADH or lNADPH. The Km value for NADH was .apprx.five times larger than the Km value for NADPH. It is inhibited by methimazole and not affected by CO. It is not induced by TTA.

AB [1-14C]Dodecylthioacetic acid (DTA), a 3-thia fatty acid, is .omega. (.omega.-1)-hydroxylated and sulfur oxygenated at about equal rates in rat liver microsomes. In prolonged incubations DTA is converted to .omega.-hydroxydodecylsulfoxyacetic acid. .omega.-Hydroxylation of DTA is catalyzed by cytochrome P450IVA1 (or a very closely related isoenzyme in the same gene family), the fatty acid .omega.-hydroxylating enzyme. It is absolutely dependent on NADPH and inhibited by CO, and lauric acid is a competing substrate. .omega.-Hydroxylation of DTA is increased by feeding tetradecylthioacetic acid (TTA), a 3-thia fatty acid, for 4 days to rats. .omega.-Hydroxylation of [1-14C] lauric acid is also induced by TTA and other 3-thia **carboxylic acids**. A close relationship was obsd. between induction of microsomal .omega.-hydroxylation of fatty acid and palmitoyl-CoA hydrolase activity. DTA is .omega.-hydroxylated at about the same rate as the physiol. substrate lauric acid. The sulfur oxygenation of DTA is catalyzed by liver microsomal **flavin-contg. monooxygenase** (FMO) (EC 1.14.13.8). It is dependent on either NADH or lNADPH. The Km value for NADH was .apprx.five times larger than the Km value for NADPH. It is inhibited by methimazole and not affected by CO. It is not induced by TTA.

L11 ANSWER 3 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1985:61628 HCAPLUS

DOCUMENT NUMBER: 102:61628

TITLE: Oxidative ether cleavage with p-nitroperbenzoic acid

AUTHOR(S): Schneider, Hans Joerg; Ahlhelm, Alfred; Mueller, Walter

CORPORATE SOURCE: Univ. Saarlandes, Saarbruecken, D-6600/11, Fed. Rep. Ger.

SOURCE: Chem. Ber. (1984), 117(11), 3297-302

CODEN: CHBEAM; ISSN: 0009-2940

DOCUMENT TYPE: Journal

LANGUAGE: German

AB The reaction of p-nitroperbenzoic acid in CHCl3 with Me2CH(CH2)3OMe and (Me2CHCH2CH2)2O leads by selective attack at C-H bonds in the .alpha.-position to the ether oxygen to hemiacetals, which decomp. to aldehydes and alcs., yielding **carboxylic acids**. Secondary alkoxy groups, as in ethoxycyclohexane, furnish Baeyer-Villiger oxidn. products of initially formed ketones. Kinetic measurements with substituted benzyl Me ethers show a Hammett reaction const. .rho. = -0.9, which is in accord with the obsd. relatively small discrimination between secondary and tertiary C-H bonds. The results are compared with similar hydroxylations of alkanes and with **monooxygenase**

reactions and point to oxenoid transition states. Radical reactions, as found with some **alkanes** are not obsd., which is shown by the small amts. of PhNO₂ (.1 to req. 10%) formed during the reaction. 13C-NMR shifts of several ethers and oxidn. products are reported.

AB The reaction of p-nitroperbenzoic acid in CHCl₃ with Me₂CH(CH₂)₃OMe and (Me₂CHCH₂CH₂)₂O leads by selective attack at C-H bonds in the **alpha**.-position to the ether oxygen to hemiacetals, which decomp. to aldehydes and alcs., yielding **carboxylic acids**. Secondary alkoxy groups, as in ethoxycyclohexane, furnish Baeyer-Villiger oxidn. products of initially formed ketones. Kinetic measurements with substituted benzyl Me ethers show a Hammett reaction const. $\rho = -0.9$, which is in accord with the obsd. relatively small discrimination between secondary and tertiary C-H bonds. The results are compared with similar **hydroxylations of alkanes** and with **monooxygenase** reactions and point to oxenoid transition states. Radical reactions, as found with some **alkanes** are not obsd., which is shown by the small amts. of PhNO₂ (.1 to req. 10%) formed during the reaction. 13C-NMR shifts of several ethers and oxidn. products are reported.

L11 ANSWER 4 OF 4 HCAPLUS COPYRIGHT 2001 ACS

ACCESSION NUMBER: 1977:498893 HCAPLUS

DOCUMENT NUMBER: 87:98893

TITLE: Biosynthesis of cutin. **.omega**.-Hydroxylation of fatty

acids by a microsomal preparation from germinating Vicia faba

AUTHOR(S): Soliday, Charles L.; Kolattukudy, P. E.

CORPORATE SOURCE: Dep. Agric. Chem., Washington State Univ., Pullman, Wash., USA

SOURCE: Plant Physiol. (1977), 59(6), 1116-21

CODEN: PLPHAY

DOCUMENT TYPE: Journal

LANGUAGE: English

AB **.omega**.-Hydroxylation of fatty acids, which is a key reaction in the biosynthesis of cutin and suberin, was demonstrated in a cell-free prepn. from a higher plant. A crude microsomal fraction (105,000-g pellet) from germinating embryonic shoots of V. faba catalyzed the conversion of palmitic acid to **.omega**.-hydroxypalmitic acid. As the crude cell-free prepn. also catalyzes the formation of other hydroxy acids

such as **.alpha**.- and **.beta**.-hydroxy acids, the **.omega** **.-hydroxylation** product was identified by gas chromatog. on a polyester column and reverse phase, high performance liq. chromatog., 2 techniques which were shown to resolve the positional isomers. Gas chromatog. anal. of the **dicarboxylic acid** obtained by CrO₃ oxidn. of the enzymic product also confirmed the identity of the enzymic **.omega**.-hydroxylation product. This enzymic **hydroxylation** required O and NADPH, but substitution of NADH resulted in nearly half the reaction rate obtained with NADPH. Maximal rates of **.omega**.-hydroxylation occurred at pH 8 and the rate increased in a sigmoidal manner with increasing concns. of palmitic acid. This **.omega**.-hydroxylation was inhibited by the classical **mixed function oxidase** inhibitors such as metal chelators (o-phenanthroline, 8-hydroxyquinoline, and **.alpha**., **.alpha**-dipyridyl), NaN₃, and thiol reagents (N-ethylmaleimide and p-chloromercuribenzoate). As expected of a hydroxylase, involving cytochrome P450, the present o-hydroxylase was inhibited by CO and this enzyme system showed unusually high sensitivity to this inhibition; 10% CO caused inhibition and 30% CO completely inhibited the reaction. Another unusual feature was that the

inhibition caused by any level of CO could not be reversed by light (420-460 nm).

AB **.omega.-Hydroxylation** of fatty acids, which is a key reaction in the biosynthesis of cutin and suberin, was demonstrated in a cell-free prepn. from a higher plant. A crude microsomal fraction (105,000-g pellet) from germinating embryonic shoots of *V. faba* catalyzed the conversion of palmitic acid to **.omega.-hydroxypalmitic acid**. As the crude cell-free prepn. also catalyzes the formation of other hydroxy acids

such as **.alpha.-** and **.beta.-hydroxy acids**, the **.omega.-hydroxylation** product was identified by gas chromatog. on a polyester column and reverse phase, high performance liq. chromatog., 2 techniques which were shown to resolve the positional isomers. Gas chromatog. anal. of the **dicarboxylic acid** obtained by CrO₃ oxidn. of the enzymic product also confirmed the identity of the enzymic **.omega.-hydroxylation** product. This enzymic **hydroxylation** required O and NADPH, but substitution of NADH resulted in nearly half the reaction rate obtained with NADPH. Maximal rates of **.omega.-hydroxylation** occurred at pH 8 and the rate increased in a sigmoidal manner with increasing concns. of palmitic acid. This **.omega.-hydroxylation** was inhibited by the classical **mixed function oxidase** inhibitors such as metal chelators (o-phenanthroline, 8-hydroxyquinoline, and **.alpha.,.alpha.-dipyridyl**), NaN₃, and thiol reagents (N-ethylmaleimide and p-chloromercuribenzoate). As expected of a hydroxylase, involving cytochrome P450, the present o-hydroxylase was inhibited by CO and this enzyme system showed unusually high sensitivity to this inhibition; 10% CO caused inhibition and 30% CO completely inhibited the reaction. Another unusual feature was that the inhibition caused by any level of CO could not be reversed by light (420-460 nm).